




ASTRONOMIA


Relativitat: Escenaris Astrofísics I: La Galàxia

La Nau Gran

Manel Perucho



<http://adc.gsfc.nasa.gov/mw>



Multiwavelength Milky Way

NATIONAL
GEOGRAPHIC




A TURBULENT HEART

4 A graph based on a radio survey reveals the whirlpool motion of molecular gas in the inner part of our galaxy: gas moving away from Earth (top half) and toward Earth (bottom half). The densest gas appears where the two dots meet.

5 Massive amounts of energy are released near the center of the Milky Way, producing

electrons that race along magnetic field lines, illuminating remnants of stellar explosions.

6 Probing even deeper into the core, a radio image details a spiral of hot gas that is falling toward what may be a black hole some 2.5 million times as massive as the sun.



a small, hot core, which will cool and fade over eons to come. Its stellar wind, streams of charged particles, rushes outward in opposite directions, like exhaust from back-to-back jet engines. This bipolarism, revealed by the Hubble Space Telescope, is common among planetary nebulae. Ultraviolet light from the star heats M2-9's gases and makes them glow.

Other types of nebulae exist in our

cloud sea. Have a look at the

8 The Hubble Space Telescope's plus sign is packed into a ring of stars, a planetary nebula. The Omega Cloud (right) is a ring of gas and dust, a type of nebula that is common in our galaxy.

Way, United States.

our 4.6-billion-year-old sun.

Peering between dust clouds toward the central bulge of the Milky Way, the Hubble Space Telescope focused on a rare, clear region in the Sagittarius star cloud (above right). These Sagittarius stars formed at different times; most are older

scope in Chile and mapped in infrared light (right). A star 20 times as massive as the sun and 10,000 times brighter, it sports a disk of circumstellar dust, shown here in false color, about 20,000 times wider than Earth's

As the sun dies, they become factories for interstellar dust. Celestial soot—the remnants of its red giant stage—surrounds the tiny hot central star of NGC 7027 (above right). Blown outward, the soot would obscure our view of the center of NGC 7027 were it not for this remarkable composite image in infrared and

strewn over huge regions along the central plane of the Milky Way, are not thick and smooth but seem as frothy as the head on a glass of beer. Supernova shock waves and stellar wind from




supernova remnant. At its heart lies a pulsar—a collapsed star—whirling 30 times a second.

Satellite galaxies of the Milky Way host equally remarkable celestial phenomena. In the Large Magellanic Cloud (above right), 180,000 light-years from

13 Glistening along the spiral arms of the galaxy, bright emission nebulae mark regions where new stars are forming. The Lagoon Nebula (above), about 5,000 light-years distant, is easily detected with the naked eye as a fuzzy spot in the southern constellation Sagittarius. Wide-field images show that it covers more of


the sky than does the full moon. Where there was once only a vast dark cloud, radiation from the brightest and most massive young star in the nebula, Herschel 36, heats and ionizes the gas across a wide region. Despite the brilliance of the Lagoon Nebula and similar objects like the famous Orion Nebula, such areas are usually little more than hot blisters on the flanks of giant interstellar clouds.




LAGOON NEBULA

History
to
which
in
Aur-
ing


IMAGE CREDIT: EUROPEAN SPACE AGENCY




Fermi Bubbles




Fermi Bubbles



Forats negres estel·lars








SN1006,
Xina, Japó



Nebulosa del Cranc
Resto de Supernova (China, 1054)






Restes de Supernova en el Núvol Mayor de Magallanes




Supernova de Tycho (1572)



Restes de Supernova en el Núvol Major de Magallanes



Massive stars explode




X-ray pictures of supernova
remnants from NASA's
Chandra Observatory
<http://chandra.harvard.edu/>


The progenitor of a Type Ia supernova




Two normal stars are in a binary pair.



The more massive star becomes a giant...




...which spills gas onto the secondary star, causing it to expand and become engulfed.




The secondary, lighter star and the core of the giant star spiral toward within a common envelope.



The common envelope is ejected, while the separation between the core and the secondary star decreases.




The remaining core of the giant collapses and becomes a white dwarf.



The aging companion star starts swelling, spilling gas onto the white dwarf.




The white dwarf's mass increases until it reaches a critical mass and explodes...



...causing the companion star to be ejected away.


Supernoves
tipus I




Gamma-Ray Bursts (Imaginary Picture)

gamma-rays are produced when
the jet (close to the light speed) breaks
out from the stellar envelope


Observer




A black hole, accretion disk and jet are
formed by the gravitational collapse of
the stellar core



A very massive star (more than 20 solar mass),
whose outer envelope (hydrogen and helium) has
been removed







12800 milions d'anys llum!!!!

Gamma-Ray Bursts (GRBs): The Long and Short of It

Long gamma-ray burst (>2 seconds' duration)



Short gamma-ray burst (<2 seconds' duration)




Associats a la formació de forats negres.


Diferents tipus.


Si el nucli resultant d'una explosió supernova té entre 1.5 i $3 M_{\text{sol}}$, el resultat és un estel de neutrons.

Si el nucli restant té una massa superior a $3 M_{\text{sol}}$, l'objecte col·lapsa sobre sí mateix i trenca l'espai-temps, formant un forat negre.




A NEUTRON STAR: SURFACE and INTERIOR






ESTRELLA DE NEUTRONES



DENSIDAD MEDIA : $\rho \sim 10^{14} \text{ g/cm}^3$



Estels
de
neutrons



Isolated Neutron Star RX J185635-3754 HST • WFPC2

PRC97-32 • ST ScI OPO • September 25, 1997


F. Walter (State University of New York at Stony Brook) and NASA



- Púlsars:
 - Estels de neutrons amb alt nivell de magnetització.
 - Descobriment: 1967.




Neutron stars and black holes in our Galaxy




Relativitat General

Com detectem forats negres? De manera indirecta.

- Emissió d'alta energia.
- Altres mètodes indirectes relacionats amb l'emissió en l'entorn de l'objecte.
- Estimant el tamany de la regió emissora.







Cygnus X-1 (8000 anys-llum).



NuSTAR (X-rays)




Cygnus X-1: Forat negre de $15 M_{\text{sun}}$ + Supergegant blava




~20 sources with detected jets
(microquasars) in the galaxy
(Massi '05, Ribó '05).

Wind-jet interaction in massive X-ray binaries

Image: NASA/ESA






Estels binaris

Microquasars


GRS 1915+105

VLA


Forat negre de
 $10-18 M_{\text{sun}}$



SS433: Objecte compacte + estel.




SS433: Objecte compacte + estel.




42 dies d'observacions (2003)

SS433: Objecte compacte + estel.




The Radio–X-ray correlation





SS433

Mecanismes de producció de radiació de molt alta energia



Mecanismes de producció de radiació de molt alta energia






J.M. Paredes, J. Martí, M. Ribó, M. Massi (2000)

LS 5039

radio


MILLIARC SEC

MILLIARC SEC




DB: PW05DAFL.vti
Cycle: 5

Volume
Var: Density





Max: 7.269e+05
Min: 0.001000



user: xparedes
Wed Nov 13 13:41:02 2013


DB: PW05DAFL.vti
Cycle: 5






Binary pulsar – gravitational waves

Hulse-Taylor pulsar. Nobel prize 1993.





El període dels pulsos es fa més llarg quan els objectes s'apropen: efecte relativista.

ones gravitatòries





ones gravitatòries



ones gravitatòries



LIGO (USA)





LISA
(delayed)




VIRGO (Italia)

ones gravitatòries




El centre de la Galàxia



Mirant al Sud, a la constel·lació de Sagitari.

El centre de la Galàxia




Mirant al Sud, a la constel·lació de Sagitari.

Wide-Field Radio Image of the Galactic Center

$\lambda = 90 \text{ cm}$

(Kassim, LaRosa, Lazio, & Hyman 1999)



El centre de la Galàxia en radio freqüències.


Infraroig



Galactic Center Survey
Hubble Space Telescope NICMOS • Spitzer Space Telescope IRAC

NASA, ESA, Q.D. Wang (University of Massachusetts, Amherst), and S. Stolovy (Caltech)

STScI-PRC09-02a




Dust in the Center of the Milky Way Galaxy

NASA / JPL-Caltech / S. Stolovy (Spitzer Science Center/Caltech)

Spitzer Space Telescope • IRAC

ssc2006-02b




Young Stars Forming in the Galactic Center


NASA / JPL-Caltech / S. V. Ramírez (NExScI/Caltech)

Spitzer Space Telescope • IRAC • MIPS • IRS

ssc2009-13a




Raigs X - XMM





Raigs gamma


HESS



1992.0









Right Ascension difference from 17h 45m 40.045s

+0.5" +0.4" +0.3" +0.2" +0.1" 0.0" -0.1" -0.2"



Relativitat General



El centre de la galàxia

Fermi Bubbles

