

Conjuntos super débilmente compactos y los espacios que generan

M. Raja (Murcia)

XIII Encuentro de la Red de Análisis Funcional y Aplicaciones

Research partially supported by



Super-reflexivity


Super-reflexivity was introduced by James as a way to isolate the isomorphic properties shared by uniformly convex or uniformly smooth Banach spaces. It turned out that the spaces that can be renormed to be uniformly convex or uniformly smooth are exactly the super-reflexive spaces (Enflo).

Super-reflexivity

Super-reflexivity was introduced by James as a way to isolate the isomorphic properties shared by uniformly convex or uniformly smooth Banach spaces. It turned out that the spaces that can be renormed to be uniformly convex or uniformly smooth are exactly the super-reflexive spaces (Enflo). The uniformly convex renorming implies that the ball (and any convex bounded set) can be peeled off in finitely many steps by successive removing of slices of small diameter.

Super-reflexivity

Super-reflexivity was introduced by James as a way to isolate the isomorphic properties shared by uniformly convex or uniformly smooth Banach spaces. It turned out that the spaces that can be renormed to be uniformly convex or uniformly smooth are exactly the super-reflexive spaces (Enflo). The uniformly convex renorming implies that the ball (and any convex bounded set) can be peeled off in finitely many steps by successive removing of slices of small diameter.



Super weakly compact operators

Beauzamy (1976) extended the notion of superreflexivity to operators, firstly under the name of “opérateurs uniformément convexifiants”.

Super weakly compact operators

Beauzamy (1976) extended the notion of superreflexivity to operators, firstly under the name of “opérateurs uniformément convexifiants”. An operator $T : Z \rightarrow X$ is *uniformly convex* if given $\varepsilon > 0$ there is $\delta > 0$ such that $\|T(x) - T(y)\| < \varepsilon$ whenever $\|x\| = \|y\| = 1$ and $\|x + y\| > 2 - \delta$.

Super weakly compact operators

Beauzamy (1976) extended the notion of superreflexivity to operators, firstly under the name of “opérateurs uniformément convexifiants”. An operator $T : Z \rightarrow X$ is *uniformly convex* if given $\varepsilon > 0$ there is $\delta > 0$ such that $\|T(x) - T(y)\| < \varepsilon$ whenever $\|x\| = \|y\| = 1$ and $\|x + y\| > 2 - \delta$. An operator $T : Z \rightarrow X$ is said *uniformly convexifying* if it becomes uniformly convex after a suitable renorming of Z .

Super weakly compact operators

Beauzamy (1976) extended the notion of superreflexivity to operators, firstly under the name of “opérateurs uniformément convexifiants”. An operator $T : Z \rightarrow X$ is *uniformly convex* if given $\varepsilon > 0$ there is $\delta > 0$ such that $\|T(x) - T(y)\| < \varepsilon$ whenever $\|x\| = \|y\| = 1$ and $\|x + y\| > 2 - \delta$. An operator $T : Z \rightarrow X$ is said *uniformly convexifying* if it becomes uniformly convex after a suitable renorming of Z .

As in the Banach space case, this property can be formulated as a “super-property”, so nowadays we speak of the class of *super weakly compact operators*, denoted \mathfrak{W}^{super} .

Super weakly compact operators

Beauzamy (1976) extended the notion of superreflexivity to operators, firstly under the name of “opérateurs uniformément convexifiants”. An operator $T : Z \rightarrow X$ is *uniformly convex* if given $\varepsilon > 0$ there is $\delta > 0$ such that $\|T(x) - T(y)\| < \varepsilon$ whenever $\|x\| = \|y\| = 1$ and $\|x + y\| > 2 - \delta$. An operator $T : Z \rightarrow X$ is said *uniformly convexifying* if it becomes uniformly convex after a suitable renorming of Z .

As in the Banach space case, this property can be formulated as a “super-property”, so nowadays we speak of the class of *super weakly compact operators*, denoted \mathfrak{W}^{super} .

It turns out that \mathfrak{W}^{super} is a symmetric closed ideal such that

$$\mathfrak{K} \subsetneq \mathfrak{W}^{super} \subsetneq \mathfrak{W}$$

Super weakly compact operators

Beauzamy (1976) extended the notion of superreflexivity to operators, firstly under the name of “opérateurs uniformément convexifiants”. An operator $T : Z \rightarrow X$ is *uniformly convex* if given $\varepsilon > 0$ there is $\delta > 0$ such that $\|T(x) - T(y)\| < \varepsilon$ whenever $\|x\| = \|y\| = 1$ and $\|x + y\| > 2 - \delta$. An operator $T : Z \rightarrow X$ is said *uniformly convexifying* if it becomes uniformly convex after a suitable renorming of Z .

As in the Banach space case, this property can be formulated as a “super-property”, so nowadays we speak of the class of *super weakly compact operators*, denoted \mathfrak{W}^{super} .

It turns out that \mathfrak{W}^{super} is a symmetric closed ideal such that

$$\mathfrak{K} \subsetneq \mathfrak{W}^{super} \subsetneq \mathfrak{W}$$

Besides, the ideal \mathfrak{W}^{super} lacks the factorisation property.

Super weakly compact sets

Super weakly compact sets

The slicing property of sets in super-reflexive spaces motivates us to introduce in 2008 the notion of *finitely dentable set*.

Super weakly compact sets

The slicing property of sets in super-reflexive spaces motivates us to introduce in 2008 the notion of *finitely dentable set*. Independently, Cheng et al. introduced in 2010 the notion of *super weakly compact set* that is equivalent to the previous one for closed convex bounded sets.

Super weakly compact sets

The slicing property of sets in super-reflexive spaces motivates us to introduce in 2008 the notion of *finitely dentable set*. Independently, Cheng et al. introduced in 2010 the notion of *super weakly compact set* that is equivalent to the previous one for closed convex bounded sets. Let us give the definition in the most simple way:

Definition

A subset $K \subset X$ is said super weakly compact if $K^{\mathcal{U}}$ is a weakly compact subset of $X^{\mathcal{U}}$ for any free ultrafilter \mathcal{U} .

Super weakly compact sets

The slicing property of sets in super-reflexive spaces motivates us to introduce in 2008 the notion of *finitely dentable set*. Independently, Cheng et al. introduced in 2010 the notion of *super weakly compact set* that is equivalent to the previous one for closed convex bounded sets. Let us give the definition in the most simple way:

Definition

A subset $K \subset X$ is said super weakly compact if $K^{\mathcal{U}}$ is a weakly compact subset of $X^{\mathcal{U}}$ for any free ultrafilter \mathcal{U} .

The relation of equivalence in the ultrapower is $(x_i)_{i \in I} \sim (y_i)_{i \in I}$ if and only if $\lim_{i \in \mathcal{U}} \|x_i - y_i\| = 0$. Note that it is enough to consider free ultrafilters on \mathbb{N} since weakly compactness is separably determined after the Eberlein-Šmul'yan theorem.

Convex super weakly compact sets

Our results combined with those from Cheng et al. give the following.

Convex super weakly compact sets

Our results combined with those from Cheng et al. give the following.

Proposition. Let X be a Banach space and $K \subset X$ a bounded closed convex subset. The following conditions are equivalent:

- (i) K is super weakly compact;
- (ii) K is finitely dentable;

Convex super weakly compact sets

Our results combined with those from Cheng et al. give the following.

Proposition. Let X be a Banach space and $K \subset X$ a bounded closed convex subset. The following conditions are equivalent:

- (i) K is super weakly compact;
- (ii) K is finitely dentable;
- (iii) K does not have the finite tree property;

Convex super weakly compact sets

Our results combined with those from Cheng et al. give the following.

Proposition. Let X be a Banach space and $K \subset X$ a bounded closed convex subset. The following conditions are equivalent:

- (i) K is super weakly compact;
- (ii) K is finitely dentable;
- (iii) K does not have the finite tree property;
- (iv) there is a reflexive Banach space Z and a super weakly compact operator $T : Z \rightarrow X$ such that $K \subset T(B_Z)$;

Convex super weakly compact sets

Our results combined with those from Cheng et al. give the following.

Proposition. Let X be a Banach space and $K \subset X$ a bounded closed convex subset. The following conditions are equivalent:

- (i) K is super weakly compact;
- (ii) K is finitely dentable;
- (iii) K does not have the finite tree property;
- (iv) there is a reflexive Banach space Z and a super weakly compact operator $T : Z \rightarrow X$ such that $K \subset T(B_Z)$;
- (v) there is a bounded uniformly convex function $f : K \rightarrow \mathbb{R}$;

Convex super weakly compact sets

Our results combined with those from Cheng et al. give the following.

Proposition. Let X be a Banach space and $K \subset X$ a bounded closed convex subset. The following conditions are equivalent:

- (i) K is super weakly compact;
- (ii) K is finitely dentable;
- (iii) K does not have the finite tree property;
- (iv) there is a reflexive Banach space Z and a super weakly compact operator $T : Z \rightarrow X$ such that $K \subset T(B_Z)$;
- (v) there is a bounded uniformly convex function $f : K \rightarrow \mathbb{R}$;
- (vi) there is an equivalent norm $\|\cdot\|$ on X which is uniformly convex on K .

Some properties of super weakly compact sets

Among the interesting properties enjoyed by the class of super weakly compact convex sets we have:

- ① the class behave quite well by usual operations: convex closed subsets, finite products, algebraic sums, linear continuous images. . . of sets from the class remain in the class;

Some properties of super weakly compact sets

Among the interesting properties enjoyed by the class of super weakly compact convex sets we have:

- 1 the class behave quite well by usual operations: convex closed subsets, finite products, algebraic sums, linear continuous images. . . of sets from the class remain in the class;
- 2 they are *uniformly Eberlein compacts*, that is, homeomorphic in the weak topology to weakly compact subsets of Hilbert spaces;

Some properties of super weakly compact sets

Among the interesting properties enjoyed by the class of super weakly compact convex sets we have:

- 1 the class behave quite well by usual operations: convex closed subsets, finite products, algebraic sums, linear continuous images. . . of sets from the class remain in the class;
- 2 they are *uniformly Eberlein compacts*, that is, homeomorphic in the weak topology to weakly compact subsets of Hilbert spaces;
- 3 they have the *Banach-Saks property*;

Some properties of super weakly compact sets

Among the interesting properties enjoyed by the class of super weakly compact convex sets we have:

- ① the class behave quite well by usual operations: convex closed subsets, finite products, algebraic sums, linear continuous images. . . of sets from the class remain in the class;
- ② they are *uniformly Eberlein compacts*, that is, homeomorphic in the weak topology to weakly compact subsets of Hilbert spaces;
- ③ they have the *Banach-Saks property*;
- ④ they have *normal structure* (and so the fixed point property for non-expansive maps) after a renorming of the ambient space;

Some properties of super weakly compact sets

Among the interesting properties enjoyed by the class of super weakly compact convex sets we have:

- 1 the class behave quite well by usual operations: convex closed subsets, finite products, algebraic sums, linear continuous images. . . of sets from the class remain in the class;
- 2 they are *uniformly Eberlein compacts*, that is, homeomorphic in the weak topology to weakly compact subsets of Hilbert spaces;
- 3 they have the *Banach-Saks property*;
- 4 they have *normal structure* (and so the fixed point property for non-expansive maps) after a renorming of the ambient space;
- 5 of course, a Banach space is super-reflexive if and only if its unit ball is super weakly compact, and a operator $T : Z \rightarrow X$ is super weakly compact if and only if $\overline{T(B_Z)}$ is super weakly compact.

Spaces generated by ideals, sets or... other spaces

Now we will consider several sorts of generation of spaces related to super-reflexivity and super weakly compactness. Let us recall the following consequence of the celebrated result of Davis, Figiel, Johnson and Pełczyński.

Spaces generated by ideals, sets or ... other spaces

Now we will consider several sorts of generation of spaces related to super-reflexivity and super weakly compactness. Let us recall the following consequence of the celebrated result of Davis, Figiel, Johnson and Pełczyński.

Theorem. For a Banach space X the following are equivalent:

- (a) There is $K \subset X$ weakly compact with $\overline{\text{span}}(K) = X$;

Spaces generated by ideals, sets or ... other spaces

Now we will consider several sorts of generation of spaces related to super-reflexivity and super weakly compactness. Let us recall the following consequence of the celebrated result of Davis, Figiel, Johnson and Pełczyński.

Theorem. For a Banach space X the following are equivalent:

- (a) There is $K \subset X$ weakly compact with $\overline{\text{span}}(K) = X$;
- (b) there are Z and $T \in \mathfrak{M}(Z, X)$ such that $\overline{T(Z)} = X$;

Spaces generated by ideals, sets or ... other spaces

Now we will consider several sorts of generation of spaces related to super-reflexivity and super weakly compactness. Let us recall the following consequence of the celebrated result of Davis, Figiel, Johnson and Pełczyński.

Theorem. For a Banach space X the following are equivalent:

- (a) There is $K \subset X$ weakly compact with $\overline{\text{span}}(K) = X$;
- (b) there are Z and $T \in \mathfrak{M}(Z, X)$ such that $\overline{T(Z)} = X$;
- (c) there are a reflexive space Z and $T \in \mathcal{L}(Z, X)$ such that $\overline{T(Z)} = X$.

Spaces generated by ideals, sets or ... other spaces

Now we will consider several sorts of generation of spaces related to super-reflexivity and super weakly compactness. Let us recall the following consequence of the celebrated result of Davis, Figiel, Johnson and Pełczyński.

Theorem. For a Banach space X the following are equivalent:

- (a) There is $K \subset X$ weakly compact with $\overline{\text{span}}(K) = X$;
- (b) there are Z and $T \in \mathfrak{M}(Z, X)$ such that $\overline{T(Z)} = X$;
- (c) there are a reflexive space Z and $T \in \mathcal{L}(Z, X)$ such that $\overline{T(Z)} = X$.

The spaces enjoying those properties are called *weakly compactly generated* and the class is denoted WCG.

Super weakly compactly generated Banach spaces

Let us consider a special sort of WCG spaces.

Super weakly compactly generated Banach spaces

Let us consider a special sort of WCG spaces.

Definition

We say that a Banach space X is super weakly compactly generated (super WCG for short) if there are Z and $T \in \mathfrak{W}^{super}(Z, X)$ such that $X = \overline{T(Z)}$.

Super weakly compactly generated Banach spaces

Let us consider a special sort of WCG spaces.

Definition

We say that a Banach space X is super weakly compactly generated (super WCG for short) if there are Z and $T \in \mathfrak{W}^{super}(Z, X)$ such that $X = \overline{T(Z)}$.

Thanks to the interpolation result shown before, X is super WCG if and only if it contains a convex super weakly compact subset with dense linear span.

Super weakly compactly generated Banach spaces

Let us consider a special sort of WCG spaces.

Definition

We say that a Banach space X is super weakly compactly generated (super WCG for short) if there are Z and $T \in \mathfrak{W}^{super}(Z, X)$ such that $X = \overline{T(Z)}$.

Thanks to the interpolation result shown before, X is super WCG if and only if it contains a convex super weakly compact subset with dense linear span. That means, “ideal-generation” and “subset-generation” notions coincide for super weak compactness.

Super weakly compactly generated Banach spaces

Let us consider a special sort of WCG spaces.

Definition

We say that a Banach space X is super weakly compactly generated (super WCG for short) if there are Z and $T \in \mathfrak{W}^{super}(Z, X)$ such that $X = \overline{T(Z)}$.

Thanks to the interpolation result shown before, X is super WCG if and only if it contains a convex super weakly compact subset with dense linear span. That means, “ideal-generation” and “subset-generation” notions coincide for super weak compactness.

However, “space-generation” is more restrictive here as we will see.

Some geometric properties of norms

The following definitions are needed...

Some geometric properties of norms

The following definitions are needed...

Given a Banach space $(X, \|\cdot\|)$:

- 1 The norm is uniformly Gâteaux (UG) smooth if for every $h \in X$

$$\sup\{\|x + th\| + \|x - th\| - 2 : x \in S_X\} = o(t) \text{ when } t \rightarrow 0.$$

Some geometric properties of norms

The following definitions are needed...

Given a Banach space $(X, \|\cdot\|)$:

- 1 The norm is uniformly Gâteaux (UG) smooth if for every $h \in X$

$$\sup\{\|x + th\| + \|x - th\| - 2 : x \in S_X\} = o(t) \text{ when } t \rightarrow 0.$$

- 2 Given a bounded set $H \subset X$, the norm is H -UG smooth if

$$\sup\{\|x + th\| + \|x - th\| - 2 : x \in S_X, h \in H\} = o(t) \text{ when } t \rightarrow 0.$$

Some geometric properties of norms

The following definitions are needed...

Given a Banach space $(X, \|\cdot\|)$:

- ① The norm is uniformly Gâteaux (UG) smooth if for every $h \in X$

$$\sup\{\|x + th\| + \|x - th\| - 2 : x \in S_X\} = o(t) \text{ when } t \rightarrow 0.$$

- ② Given a bounded set $H \subset X$, the norm is H -UG smooth if

$$\sup\{\|x + th\| + \|x - th\| - 2 : x \in S_X, h \in H\} = o(t) \text{ when } t \rightarrow 0.$$

- ③ The norm is uniformly Fréchet smooth if it is B_X -UG.

Some geometric properties of norms

The following definitions are needed...

Given a Banach space $(X, \|\cdot\|)$:

- 1 The norm is uniformly Gâteaux (UG) smooth if for every $h \in X$

$$\sup\{\|x + th\| + \|x - th\| - 2 : x \in S_X\} = o(t) \text{ when } t \rightarrow 0.$$

- 2 Given a bounded set $H \subset X$, the norm is H -UG smooth if

$$\sup\{\|x + th\| + \|x - th\| - 2 : x \in S_X, h \in H\} = o(t) \text{ when } t \rightarrow 0.$$

- 3 The norm is uniformly Fréchet smooth if it is B_X -UG.

- 4 The norm is strongly UG smooth if it is H -UG smooth for some bounded and linearly dense subset $H \subset X$.

Generation by several kind of spaces

The relationships among different kinds of generated Banach spaces were clearly exposed by Fabian, Godefroy, Hájek and Zizler in the following result of 2003.

Theorem. For a Banach space X consider the assertions:

- (i) X is Hilbert-generated.

Generation by several kind of spaces

The relationships among different kinds of generated Banach spaces were clearly exposed by Fabian, Godefroy, Hájek and Zizler in the following result of 2003.

Theorem. For a Banach space X consider the assertions:

- (i) X is Hilbert-generated.
- (ii) X is superreflexive-generated.

Generation by several kind of spaces

The relationships among different kinds of generated Banach spaces were clearly exposed by Fabian, Godefroy, Hájek and Zizler in the following result of 2003.

Theorem. For a Banach space X consider the assertions:

- (i) X is Hilbert-generated.
- (ii) X is superreflexive-generated.
- (iii) X is generated by the ℓ_2 -sum of superreflexive spaces.

Generation by several kind of spaces

The relationships among different kinds of generated Banach spaces were clearly exposed by Fabian, Godefroy, Hájek and Zizler in the following result of 2003.

Theorem. For a Banach space X consider the assertions:

- (i) X is Hilbert-generated.
- (ii) X is superreflexive-generated.
- (iii) X is generated by the ℓ_2 -sum of superreflexive spaces.
- (iv) X admits an equivalent strongly UG smooth norm.

Generation by several kind of spaces

The relationships among different kinds of generated Banach spaces were clearly exposed by Fabian, Godefroy, Hájek and Zizler in the following result of 2003.

Theorem. For a Banach space X consider the assertions:

- (i) X is Hilbert-generated.
- (ii) X is superreflexive-generated.
- (iii) X is generated by the ℓ_2 -sum of superreflexive spaces.
- (iv) X admits an equivalent strongly UG smooth norm.
- (v) X is WCG and admits an equivalent UG smooth norm.

Generation by several kind of spaces

The relationships among different kinds of generated Banach spaces were clearly exposed by Fabian, Godefroy, Hájek and Zizler in the following result of 2003.

Theorem. For a Banach space X consider the assertions:

- (i) X is Hilbert-generated.
- (ii) X is superreflexive-generated.
- (iii) X is generated by the ℓ_2 -sum of superreflexive spaces.
- (iv) X admits an equivalent strongly UG smooth norm.
- (v) X is WCG and admits an equivalent UG smooth norm.
- (vi) X is a subspace of a Hilbert-generated space.

Generation by several kind of spaces

The relationships among different kinds of generated Banach spaces were clearly exposed by Fabian, Godefroy, Hájek and Zizler in the following result of 2003.

Theorem. For a Banach space X consider the assertions:

- (i) X is Hilbert-generated.
- (ii) X is superreflexive-generated.
- (iii) X is generated by the ℓ_2 -sum of superreflexive spaces.
- (iv) X admits an equivalent strongly UG smooth norm.
- (v) X is WCG and admits an equivalent UG smooth norm.
- (vi) X is a subspace of a Hilbert-generated space.

Then $(i) \Rightarrow (ii) \Rightarrow (iii) \Rightarrow (iv) \Rightarrow (v) \Rightarrow (vi)$.

Generation by several kind of spaces

The relationships among different kinds of generated Banach spaces were clearly exposed by Fabian, Godefroy, Hájek and Zizler in the following result of 2003.

Theorem. For a Banach space X consider the assertions:

- (i) X is Hilbert-generated.
- (ii) X is superreflexive-generated.
- (iii) X is generated by the ℓ_2 -sum of superreflexive spaces.
- (iv) X admits an equivalent strongly UG smooth norm.
- (v) X is WCG and admits an equivalent UG smooth norm.
- (vi) X is a subspace of a Hilbert-generated space.

Then $(i) \Rightarrow (ii) \Rightarrow (iii) \Rightarrow (iv) \Rightarrow (v) \Rightarrow (vi)$. Moreover, no one of these implications can be reversed in general.

Renorming characterisation of super WCG

Since \mathfrak{M}^{super} does not have the factorisation property, the class of spaces generated by this ideal could be larger than the class of super-reflexive generated Banach spaces.

Renorming characterisation of super WCG

Since \mathfrak{M}^{super} does not have the factorisation property, the class of spaces generated by this ideal could be larger than the class of super-reflexive generated Banach spaces. It is a natural question to know whether this class be placed among the six categories of the previous theorem?

Renorming characterisation of super WCG

Since \mathfrak{M}^{super} does not have the factorisation property, the class of spaces generated by this ideal could be larger than the class of super-reflexive generated Banach spaces. It is a natural question to know whether this class be placed among the six categories of the previous theorem?

The answer is that they are the spaces fulfilling statement “(iv)”

Renorming characterisation of super WCG

Since \mathfrak{M}^{super} does not have the factorisation property, the class of spaces generated by this ideal could be larger than the class of super-reflexive generated Banach spaces. It is a natural question to know whether this class be placed among the six categories of the previous theorem?

The answer is that they are the spaces fulfilling statement “(iv)”

Theorem

A Banach space X is super WCG if and only if it admits an equivalent strongly UG smooth norm.

Renorming characterisation of super WCG

Since \mathfrak{W}^{super} does not have the factorisation property, the class of spaces generated by this ideal could be larger than the class of super-reflexive generated Banach spaces. It is a natural question to know whether this class be placed among the six categories of the previous theorem?

The answer is that they are the spaces fulfilling statement “(iv)”

Theorem

A Banach space X is super WCG if and only if it admits an equivalent strongly UG smooth norm.

Our proof strongly depends on the symmetry of the ideal \mathfrak{W}^{super} .

Strong generation

Sometimes the linear density of a nice generating set is not enough to transfer good properties to the space. We require one more definition:

Strong generation

Sometimes the linear density of a nice generating set is not enough to transfer good properties to the space. We require one more definition:

A Banach space X is said to be *strongly generated by a subset* $K \subset X$ if for any weakly compact $H \subset X$ and $\varepsilon > 0$ there is $n \in \mathbb{N}$ such that $H \subset nK + \varepsilon B_X$.

Strong generation

Sometimes the linear density of a nice generating set is not enough to transfer good properties to the space. We require one more definition:

A Banach space X is said to be *strongly generated by a subset* $K \subset X$ if for any weakly compact $H \subset X$ and $\varepsilon > 0$ there is $n \in \mathbb{N}$ such that $H \subset nK + \varepsilon B_X$.

Obviously, the definition admits variations for “space-generation” and “ideal-generation”.

Strong generation

Sometimes the linear density of a nice generating set is not enough to transfer good properties to the space. We require one more definition:

A Banach space X is said to be *strongly generated by a subset* $K \subset X$ if for any weakly compact $H \subset X$ and $\varepsilon > 0$ there is $n \in \mathbb{N}$ such that $H \subset nK + \varepsilon B_X$.

Obviously, the definition admits variations for “space-generation” and “ideal-generation”.

Let us mention here that several good properties of the space $L_1(\mu)$ for μ finite can be understood from the fact that it is strongly Hilbert generated.

Strong generation and renorming

The class of Banach spaces which are strongly generated by super weakly compact convex sets (denoted S^2WCG) have a remarkable renorming property.

Strong generation and renorming

The class of Banach spaces which are strongly generated by super weakly compact convex sets (denoted S^2WCG) have a remarkable renorming property.

Theorem

Let X be a S^2WCG Banach space. Then there is an equivalent norm on X such that its restriction to any reflexive subspace is both uniformly convex and uniformly Fréchet smooth.

Strong generation and renorming

The class of Banach spaces which are strongly generated by super weakly compact convex sets (denoted S^2WCG) have a remarkable renorming property.

Theorem

Let X be a S^2WCG Banach space. Then there is an equivalent norm on X such that its restriction to any reflexive subspace is both uniformly convex and uniformly Fréchet smooth.

This result generalises the known result for some super-reflexive generated Banach spaces like Lebesgue-Bochner spaces $L_1(\mu, X)$ with X super-reflexive (Fabian-Lajara).

Strong generation and renorming

The class of Banach spaces which are strongly generated by super weakly compact convex sets (denoted S^2WCG) have a remarkable renorming property.

Theorem

Let X be a S^2WCG Banach space. Then there is an equivalent norm on X such that its restriction to any reflexive subspace is both uniformly convex and uniformly Fréchet smooth.

This result generalises the known result for some super-reflexive generated Banach spaces like Lebesgue-Bochner spaces $L_1(\mu, X)$ with X super-reflexive (Fabian-Lajara).


On the other hand, there are S^2WCG spaces which are not super-reflexively generated.

References

- B. BEAUZAMY, Operateurs uniformement convexifiants, *Studia Math.* 57 (2) (1976) 103–139.
- L. CHENG, Q. CHENG, B. WANG, W. ZHANG, On super-weakly compact sets and uniformly convexifiable sets, *Studia Math.* 199 (2) (2010) 145–169.
- M. FABIAN, G. GODEFROY, P. HAJEK, V. ZIZLER, Hilbert-generated spaces, *J. Funct. Anal.* 200 (2) (2003) 301–323.
- M. FABIAN, S. LAJARA, Smooth renormings of the LebesgueBochner function space $L_1(\mu, X)$, *Studia Math.* 209 (3) (2012) 247–265.
- M. RAJA, Finitely dentable functions, operators and sets. *J. Convex Anal.* 15 (2008), no. 2, 219–233.
- M. RAJA, Super WCG Banach spaces. *J. Math. Appl.* 439 (2016), no. 1, 183–196.

Announcement

NonLinear Functional Analysis



[Start](#) [Main speakers](#) [Short talks](#) [Schedule](#) [Registration](#) [Sending abstract](#) [Venue](#) [Committee](#) [Sponsors](#)

CONFERENCE IN NON LINEAR FUNCTIONAL ANALYSIS

The Conference on Non-Linear Functional Analysis, will be held at the Universitat Politècnica de Valencia (Spain) on **17-20 October 2017**. This is a joint project of the Universitat de València, Universidad de Murcia and Universitat Politècnica de València that includes the 5th Workshop on Functional Analysis and the Workshop on Infinite Dimensional Analysis Valencia 2017. You can find information about previous editions at <http://www.um.es/beca/workshop2016/> and <http://mate.dm.uba.ar/~widaba14/> (the first edition of this one was held in Buenos Aires).

Each day we will have two or three plenary lectures in the morning and after them the two workshops will run in parallel sessions. We intend to have a wide audience with interest in any field in Functional Analysis. Main topics of the Meeting will be Banach space theory, polynomials and multilinear mappings, vector measures and complex analysis in finite or infinite dimensional spaces, but of course contributions in any other area of Functional Analysis will be very much welcome.

You can contact us by sending an e-mail to banach@um.es

More information soon at <http://www.um.es/beca/NoLiFA/>