



remaining forests and restoring forest ecosystem resilience. However, the plans for implementing these goals are misguided.

The EU Strategy commits to “strictly protecting...primary and old-growth forests.” Known primary forests represent 0.7% of Europe’s forest area, of which only 46% are strictly protected (2). Strictly protecting the remaining forests is the right strategy, given that forest management seriously threatens remaining primeval and old forests in Europe, including the Białowieża Forest, Carpathian forests, and Scandinavian Mountains Green Belt (3, 4). However, the EU Strategy overlooks the need for spatial coherence. Tiny protected patches of forest will be of little relevance if intensive forestry prevails in the surrounding matrix. An ecosystem-based approach to forest management must complement protection efforts.

The EU Strategy also commits to restoring forest ecosystems but only offers planting more than 3 billion trees as a concrete action. Planting trees in forest habitats may have detrimental effects (5, 6). The EU Strategy doesn’t specify what kinds of trees should be planted, a necessary detail given that the forestry sector in most countries increasingly undermines ecosystem functionality by introducing exotic species believed to be better adapted to future climates (7).

Instead of planting trees, conservationists should focus on reducing the rate of forest degradation (namely, tree harvesting) and supporting natural renewal processes. The EU Strategy does not consider natural

restoration of spontaneously regenerating forests, the surface of which has decreased 7% worldwide over the past 30 years (8). Simply allowing the forests to naturally regrow often results in more trees at much lower costs than planting (6). The EU Strategy should advocate a hands-off strategy for safeguarding ecosystemic responses to degradation and environmental change. Policies should support the natural recruitment and selection for trees with greater resistance to insect attacks or extreme events (9) and biomass-rich forests with closed canopies (10), and they should prevent road construction through valuable forest patches (11). Actions aimed at promoting the adaptation of forests to global change and increasing their resilience should be based on all available ecological science and require a more complex view than a simplistic planting strategy. The EU Forest Strategy planned for 2021 should focus on developing a holistic approach with a clearly defined timetable of actions.

Nuria Selva^{1*}, Przemysław Chylarecki², Bengt-Gunnar Jonsson^{3,4}, Pierre L. Ibisch⁵

¹Institute of Nature Conservation Polish Academy of Sciences, 31-120 Kraków, Poland. ²Museum and Institute of Zoology, Polish Academy of Sciences, 00-679 Warszawa, Poland. ³Department of Natural Sciences, Mid Sweden University, SE-851 70 Sundsvall, Sweden. ⁴Department of Wildlife, Fish, and Environmental Studies, Swedish University of Agricultural Sciences, SE-901 83 Umeå, Sweden. ⁵Centre for Economics and Ecosystem Management, Eberswalde University for Sustainable Development, 16-225 Eberswalde, Germany.
*Corresponding author.
Email: nuriselve@gmail.com

REFERENCES AND NOTES

1. European Commission, “EU Biodiversity Strategy for 2030.” (2020); https://ec.europa.eu/environment/nature/biodiversity/strategy/index_en.htm.
2. F. M. Sabatini *et al.*, *Divers. Distrib.* **24**, 1426 (2018).
3. E. Stokstad, *Science* **358**, 1240 (2017).
4. B.-G. Jonsson *et al.*, *Forests* **10**, 564 (2019).
5. M. Zmihorski *et al.*, *Science* **361**, 238 (2018).
6. K. D. Holl, P. H. S. Brancalion, *Science* **368**, 580 (2020).
7. F. Krumm, L. Vítková, Eds., “Introduced tree species in European forests: opportunities and challenges” (European Forest Institute, 2016).
8. Food and Agriculture Organization of the United Nations and UN Environment Programme, *The State of the World’s Forests: Forests, Biodiversity and People* (FAO and UNEP, 2020).
9. D. L. Six, C. Vergobbi, M. Cutter, *Front. Plant Sci.* **9**, 993 (2018).
10. F. Zellweger *et al.*, *Science* **368**, 772 (2020).
11. P. L. Ibisch *et al.*, *Science* **354**, 1423 (2016).

10.1126/science.abc9892

Afforestation falls short as a biodiversity strategy

The recent EU Biodiversity Strategy for 2030 (1) recognizes the importance of biodiversity for increasing our resilience to natural disasters and pandemics and, thus, for human well-being. Although

it proposes ambitious measures such as reversing pollinator decline and controlling invasive species, it also introduces the ill-advised idea of planting 3 billion trees.

Massive tree plantation programs (2, 3) have been strongly criticized by the scientific community for their negative ecological and economic impacts and their limited role in climate change and CO₂ mitigation (4–8). The specific number of trees proposed in the EU Strategy suggests a lack of a serious, science-based ecological assessment of actual restoration needs. Meeting such a target could threaten biodiverse treeless ecosystems (4, 6, 7, 9) and would waste an opportunity to implement ecologically sound management practices to restore fully functionally integrated mosaics of natural, seminatural, and sustainable agricultural ecosystems.

Massive tree planting could also substantially change the fire regime, especially given the increasing frequency of heat waves and droughts in an area with high population density (10). The probability of large intense fires that threaten biodiversity and human assets is largely influenced by the type, amount, and continuity of biomass. Therefore, determining how many trees should be planted is less important than figuring out the most safe and effective conservation strategy.

We need to move toward optimizing our landscapes. A diverse mosaic of nature-based production systems should be interspersed with protected natural areas to maximize biodiversity, resilience, and ecosystem services. Trees are not synonymous with biodiversity. Policy-makers and society need to internalize this message to make proper decisions in the context of environmental and health crises.

Susana Gómez-González^{1,2*}, Raúl Ochoa-Hueso¹, Juli G. Pausas³

¹Departamento de Biología-IVAGRO, Universidad de Cádiz, Puerto Real, Spain. ²Center for Climate and Resilience Research (CR)², Santiago, Chile. ³Centro de Investigaciones sobre Desertificación (CIDE-CSIC), Valencia, Spain.
*Corresponding author.
Email: susana.gomez@uca.es

REFERENCES AND NOTES

1. European Commission, “EU Biodiversity Strategy for 2030” (2020); https://ec.europa.eu/environment/nature/biodiversity/strategy/index_en.htm.
2. International Union for Conservation of Nature, *The Bonn Challenge* (2020); www.bonnchallenge.org.
3. J.-F. Bastin *et al.*, *Science* **365**, 76 (2019).
4. W. J. Bond *et al.*, *Trends Ecol. Evol.* **34**, 963 (2019).
5. D. Baldocchi, J. Penuelas, *Glob. Change Biol.* **25**, 1191 (2019).
6. J. W. Veldman *et al.*, *Science* **366**, eaay7976 (2019).
7. K. D. Holl, P. H. S. Brancalion, *Science* **368**, 580 (2020).
8. M. Jiang *et al.*, *Nature* **580**, 227 (2020).
9. J. Scurlock, D. Hall, *Glob. Change Biol.* **4**, 229 (1998).
10. D. de Rigo *et al.*, “Forest fire danger extremes in Europe under climate change: Variability and uncertainty” (JRC Tech. Rep. 28926, Publications Office of the European Union, Luxembourg, 2017).

10.1126/science.abd3064