

RESEARCH ARTICLE OPEN ACCESS

Cross-Continental Analysis Shows That Disturbance Effects on Reptile Body Condition Do Not Predict Abundance Responses

Kristina J. Macdonald^{1,2}  | Don A. Driscoll²  | Michael D. Craig³  | Robert A. Davis⁴  | Steven J. Hromada⁵  | C. M. Gienger⁶  | Lee A. Fitzgerald⁷  | Daniel J. Leavitt⁸  | Danielle K. Walkup⁹  | Rickard Abom¹⁰  | Adrian F. Wayne¹¹  | Gordon R. Friend¹² | Brent Johnson¹¹ | Danielle Stokeld¹³  | Chris R. Dickman¹⁴  | Scott A. Thompson^{15,16}  | Graham G. Thompson^{3,15}  | Daniel F. Bohórquez Fandiño^{11,14} | Christopher K. Woolley¹⁷  | Annabel L. Smith¹⁸  | Calum Irvine³ | Tim S. Jessop¹⁹  | Jade E. Keehn²⁰ | Chris R. Feldman²¹  | Xavier Santos²²  | Josabel Belliure²³  | Juli G. Pausas²⁴  | Stacey L. Weiss²⁵  | Patricia A. Fleming²⁶  | Shannon Dundas²⁷  | Dylan M. Westaway²⁸  | Sabrina E. Duncan²⁹ | Tim S. Doherty^{11,14} 

¹Research Institute for the Environment and Livelihoods, Charles Darwin University, Darwin, Northern Territory, Australia | ²School of Life and Environmental Sciences, Deakin University, Melbourne, Victoria, Australia | ³School of Biological Sciences, University of Western Australia, Perth, Western Australia, Australia | ⁴School of Science, Edith Cowan University, Perth, Western Australia, Australia | ⁵Fresno Chaffee Zoo, Fresno, California, USA | ⁶Department of Biology and Center of Excellence for Field Biology, Austin Peay State University, Clarksville, Tennessee, USA | ⁷Biodiversity Research and Teaching Collections, Department of Ecology and Conservation Biology, Texas A&M University, College Station, Texas, USA | ⁸United State Fish and Wildlife Service, Falls Church, Virginia, USA | ⁹Natural Resources Institute, Texas A&M University, College Station, Texas, USA | ¹⁰College of Science and Engineering, James Cook University, Townsville, Queensland, Australia | ¹¹Biodiversity and Conservation Science, Department of Biodiversity, Conservation and Attractions, Western Australia, Australia | ¹²Department of Conservation and Land Management, Wildlife Research Centre, Perth, Western Australia, Australia | ¹³NSW Department of Climate Change, Energy, the Environment and Water, Sydney, New South Wales, Australia | ¹⁴School of Life and Environmental Sciences, The University of Sydney, Sydney, New South Wales, Australia | ¹⁵Terrestrial Ecosystems, Perth, Western Australia, Australia | ¹⁶School of Molecular and Life Sciences, Curtin University, Perth, Western Australia, Australia | ¹⁷Centre for Biodiversity and Restoration Ecology, School of Biological Sciences, Te Herenga Waka—Victoria University of Wellington, Wellington, New Zealand | ¹⁸School of the Environment, University of Queensland, Brisbane, Queensland, Australia | ¹⁹Conservation and Restoration Science Branch, Science, Economics, and Insights Division, NSW Department of Planning and Environment, Sydney, New South Wales, Australia | ²⁰Wildlife Conservation Biologist, Oregon Department of Fish and Wildlife, Central Point, Oregon, USA | ²¹Department of Biology, University of Nevada, Reno, Nevada, USA | ²²Centro de Investigação Em Biodiversidade e Recursos Genéticos da Universidade do Porto (CIBIO/InBIO), Vairão, Portugal | ²³Global Change Ecology and Evolution Research Group (GloCEE), Department of Life Sciences, University of Alcalá, Madrid, Spain | ²⁴Centro de Investigaciones Sobre Desertificación (CIDE-CSIC), Valencia, Spain | ²⁵Biology Department, University of Puget Sound, Tacoma, Washington, USA | ²⁶Environmental and Conservation Sciences, Harry Butler Institute, Murdoch University, Perth, Western Australia, Australia | ²⁷NSW Department of Primary Industries and Regional Development, Vertebrate Pest Research, Sydney, New South Wales, Australia | ²⁸Gulbali Institute, School of Agricultural, Environmental and Veterinary Sciences, Charles Sturt University, Albury, New South Wales, Australia | ²⁹Thünen Institute of Sea Fisheries, Bremerhaven, Germany

Correspondence: Kristina J. Macdonald (kristina.macdonald@cdu.edu.au)

Received: 28 January 2025 | **Revised:** 16 April 2025 | **Accepted:** 17 April 2025

Funding: This work was supported by the Australian Research Council (LP170101134, AD9031596), the Australian Research Grant Scheme (A18615230), Barrick Kanowna, Charles Darwin University, Edith Cowan University, Holsworth Wildlife Research Endowment, Minerals Research Institute of Western Australia, M.J. Murdock Charitable Trust, Australian Government's National Environmental Science Program, Northern Australian Environmental Resources Hub, Norilsk Nickel, Northern Territory Department of Lands, Planning and Environment, OMG Cawse Nickel Operations, Parks Victoria, Placer Dome Asia Pacific Kalgoorlie West Operations, and the Threatened Species Recovery Hub.

Keywords: biodiversity monitoring | fire | habitat fragmentation | invasive species | mining | physiological impacts | population decline | population fitness

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2025 The Author(s). *Global Change Biology* published by John Wiley & Sons Ltd.

ABSTRACT

Ecological disturbances are discrete events that alter or transform the physical, chemical, or biological characteristics of ecosystems. Disturbance can cause animal populations to decline and, according to the risk-disturbance hypothesis and population collapse framework, these declines can be predicted by declines in animal body condition. However, no research has empirically examined the general relationship between body condition and abundance, nor their relationship in response to disturbance. We used a combined dataset representing 33 studies and >42,000 observations of 75 species from Australia, New Zealand, Spain and the United States of America to test predictions relating to the relationship between reptile body condition and abundance. We first investigated the relationship at the site level and then used meta-analytical models to test whether populations showed linked changes in abundance and body condition in response to disturbance. We further tested whether key environmental and species traits influenced this relationship and whether there was a time-lagged effect of body condition responses on abundance. We found a positive relationship between mean reptile body condition and abundance at the site level. However, the relationship was largely lost when investigating population responses to disturbance. As such, our results provided no support for the risk-disturbance hypothesis and limited support for the population collapse framework. Therefore, the impacts of disturbance on reptile body condition cannot be assumed to reflect or predict abundance responses. We provide a new conceptual framework that shows how disturbances can modify or uncouple the relationship between abundance and body condition by influencing underlying drivers, such as predation, competition and resource availability. Monitoring programs that infer population impacts based on changes in body condition should first confirm the relationship between these two variables in the relevant study system.

1 | Introduction

Disturbances are discrete events of biotic or abiotic origin that cause either temporary or permanent changes to the physical, chemical or biological characteristics of ecosystems (Battisti et al. 2016). Natural disturbances have played important roles in the functioning and evolution of ecosystems for millions of years (Battisti et al. 2016; Pausas and Keeley 2009). More recently, though, anthropogenic disturbances have proliferated globally, including human disruption of natural disturbance regimes (IPBES 2019; Parmesan et al. 2022). Disturbances can have direct effects on wildlife, such as injury, infection or mortality, as well as indirect effects via altered environmental conditions, habitat structure, food resources and species interactions (Santos et al. 2022; Sergio et al. 2018).

Research into the effects of disturbance on wildlife has focused primarily on community and population level responses (Cordier et al. 2021; Hillebrand and Kunze 2020; Keinath et al. 2017; Quesnelle et al. 2014; Sousa 1984). Response metrics such as species richness and occupancy provide insight into the loss and gain of species following disturbance, while measures of abundance provide insight into changing population size (Deng et al. 2023; Williams et al. 2022). On the other hand, individual response metrics of animal health or fitness can encompass indirect effects of disturbance by considering the effects on individuals persisting in disturbed landscapes. As such, individual response metrics are expected to act as an early indicator of future population change (Cerini et al. 2023). For instance, lower body condition of male white-plumed honeyeaters (*Ptilotula penicillatus*) following heat exposure reduced their probability of survival in the following spring (Gardner et al. 2016).

Body condition is an individual response metric that acts as a proxy for energy stores and fitness (ability to survive and reproduce) (Labocha et al. 2014). This metric is typically calculated using the mass and size of individual animals, making it an easily accessible indicator of health compared to other alternatives (e.g., stress or immune function measures). Given the

connection between body condition and reproductive success (Lannoo and Stiles 2017; Reading 2004; Testa and Adams 1998), and body condition and mortality risk (Aubry et al. 2013; Shine et al. 2001; Stewart et al. 2021), the effects of disturbances on body condition are often thought to reflect broader population responses (e.g., Cerini et al. 2023; Cogălniceanu et al. 2021; Janin et al. 2011; Macdonald et al. 2023). However, the generalized relationship between body condition and abundance is poorly resolved (Gould et al. 2024; Macdonald et al. 2023).

The risk-disturbance hypothesis and population collapse framework predict that population declines follow reductions in individual body condition (Cerini et al. 2023; Frid and Dill 2002). The risk-disturbance hypothesis predicts that the impacts of disturbance on individuals will be characterized by changes in perceived risks, analogous to predation risk, where animals become hyper-vigilant, but at the cost of foraging and reproductive success (Frid and Dill 2002). Similarly, the population collapse framework uses individual body condition, along with other individual response metrics, to predict both concurrent and future population collapse following environmental stress (Cerini et al. 2023). Both frameworks acknowledge important assumptions and caveats, but to date, little work has explicitly tested these predictions.

A positive relationship between body condition and population abundance is expected because individuals with poor body condition can have lower fitness and higher mortality, leading to population declines (e.g., Lannoo and Stiles 2017; Madsen et al. 2023; Reading 2004; Shine et al. 2001). On the other hand, a negative relationship between body condition and abundance could arise when the relationship is driven by intra-specific competition (Herrando-Pérez et al. 2012; Stewart et al. 2005). However, there are few long-term studies explicitly assessing the links between body condition, demography, and population growth trajectories (e.g., Gardner et al. 2016; Pigeon et al. 2017; Reading 2007). Of the few studies investigating the concurrent impacts of disturbances on body condition and population metrics, some have found a positive relationship (Emery et al. 2021; Keehn et al. 2019; Lunghi et al. 2018;

Smith 2018), no relationship, or a negative relationship (Gould et al. 2024; Grimm-Seyfarth et al. 2018; Keyser et al. 2005; Laidre et al. 2006; Welsh et al. 2008). Furthermore, there can be time lags, with body condition an important predictor of annual survival in the following activity period (Madsen et al. 2023). This range of responses reported in the literature emphasizes the need for further research and empirical synthesis.

Reptiles provide a particularly good model for addressing this research question; they are widespread globally, ecologically important, relatively easy to capture compared to other vertebrates (e.g., birds), and the data needed to calculate body condition are relatively simple to collect (Cox et al. 2022; Macdonald et al. 2023; Schwarzkopf 2022). Many reptiles are threatened by habitat disturbances such as fire, invasive species, drought, and habitat fragmentation (Böhm et al. 2016; Cox et al. 2022; Tingley et al. 2019). Conversely, some reptile species rely on regular disturbance to maintain suitable habitat. For example, Indigenous fire management in Australia and fire restoration in the United States of America can promote reptile diversity by creating habitat for early successional species (Fair and Henke 1997; Legge et al. 2024). Global meta-analyses have shown that disturbances generally have negative impacts on reptile body condition (Macdonald et al. 2023) and abundance (Doherty et al. 2020). However, there was a high level of unexplained variance in the results of these studies (Doherty et al. 2020; Macdonald et al. 2023) and no research has systematically investigated the relationship between body condition and abundance responses to disturbance for any taxonomic group.

Life history traits and environmental characteristics can be important mediators of wildlife responses to disturbance (Sergio et al. 2018), including effects on abundance and body condition. Disturbance characteristics, such as disturbance type and time since disturbance, could influence how carrying capacity is affected and, in turn, determine whether abundance and body condition show similar or different responses to disturbance. For example, disturbance types that result in direct mortality (e.g., invasive predators) may not influence carrying capacity, whereas disturbances that reduce habitat (e.g., logging) can reduce carrying capacity. In addition, populations exposed to press disturbances (i.e., ongoing, continuous disturbances, such as invasive species or climatic change) may be able to adapt behaviorally to regain and maintain body condition (Rodríguez-Prieto et al. 2010). Species traits could also influence behavioral responses to disturbance, which in turn could affect body condition and abundance (Böhm et al. 2016; Lazzari et al. 2022).

Examining the relationship between reptile abundance and body condition and their relationship with disturbance provides an opportunity to test the risk-disturbance hypothesis and population collapse framework (Cerini et al. 2023; Frid and Dill 2002). We combined more than 42,000 data points from 33 studies across 75 reptile species to test the following three predictions:

1. Average individual body condition will be positively associated with abundance at the site-survey level, reflecting the mechanistic relationship between body condition, survival, and reproduction (Cerini et al. 2023; Reading 2004; Shine et al. 2001).

2. Body condition responses to disturbance will be positively correlated with abundance responses to disturbance, indicating that individual fitness consequences drive population change (Cerini et al. 2023; Frid and Dill 2002). Alongside this prediction, we investigated the potential role of disturbance characteristics and species traits in influencing the relationship between the two metrics.
3. As an alternative to prediction 2, body condition responses to disturbance in one year will be positively correlated with abundance responses to disturbance in the following year due to time-lagged effects of body condition on population growth, such as changes in reproductive output (Gardner et al. 2016; Madsen et al. 2023).

2 | Methods

Between October and December 2023, suitable datasets were identified through three main methods: (1) reviewing the data sources of a recent global meta-analysis investigating reptile body condition responses to disturbance (Macdonald et al. 2023), (2) using ‘Scopus’ to identify prominent authors publishing in the field of reptile abundance responses to disturbance, and (3) asking existing networks and research collaborators for recommendations. Adopting a formal systematic review approach was not suitable for this study because it was necessary to use individual animal and site-survey level data (often unpublished) and to consult with the data custodians (author group) regarding data cleaning and interpretation.

Datasets were deemed suitable for our research questions and meta-analytical approach if they (1) systematically surveyed reptiles (e.g., live trapping, timed searches) in both disturbed and undisturbed sites and (2) recorded count data at the species level and individual snout-vent length and weight measurements.

We formed a research collaboration that brought together 33 existing datasets meeting our research requirements (see Table S1 AND S2, “Data Sources,” and Macdonald 2025 for secondary dataset details). These datasets came from studies in Australia, New Zealand, Spain, and the United States of America, which investigated 10 disturbance types across 75 reptile species from 13 families (Table S3).

2.1 | Data Processing

We used the R software for all data processing and analysis (R Core Team 2022). For the abundance data, recaptured individuals were excluded from datasets when recorded. In addition, in datasets where sampling effort varied across sites, counts were converted to capture rates by standardizing counts by a unit of survey effort. As such, capture rates were used as a proxy of relative abundance.

For body condition data, we excluded gravid and deceased individuals (when recorded) to avoid potential weight confounding. In addition, we removed suspected measurement or data entry errors. To identify likely errors, we (1) visually assessed the range of raw snout-vent and mass values against known species

information to identify clear outliers, (2) visually assessed scatterplots of mass and snout-vent length, and (3) visually assessed the distribution of Scaled Mass Index values (explained below). Individuals with greater than double or less than half the mass of individuals of a similar snout-vent length size were excluded. The final Scaled Mass Index was calculated after errors were excluded from datasets.

2.1.1 | Scaled Mass Index

The Scaled Mass Index (SMI) is a calculation of body condition that controls for the allometric relationship between mass and growth and is robust to sexual size dimorphism (Peig and Green 2010). SMI scales the mass of individuals to the mean length of the population along a population specific allometric relationship between length and mass. Therefore, SMI values are in mass units, making results easily interpretable. We used the methods from Peig and Green (2009) to calculate the SMI of each species within each dataset. We fitted standardized major axis models using the log(mass) and log(snout-vent length) of populations using the SMTR package v3.4–8 (Warton et al. 2012). We then used the slope coefficient from the model outputs as the scaling coefficient (b_{SMA}) in subsequent SMI calculations:

$$\hat{M}_i = M_i \left[\frac{L_0}{L_i} \right]^{b_{SMA}}$$

where the scaled mass of an individual ($\hat{M}_i$) is calculated using the mass of an individual (M_i), the mean snout to vent length (L_0), the snout to vent length of the individual (L_i) and the slope coefficient from the standardized major axis model (b_{SMA}).

2.2 | Site-Survey Dataset

To investigate the relationship between relative abundance and body condition, we compiled a dataset of site-survey responses (Figure 1). For each species within each study, we compiled capture rates and the mean SMI for each site-survey when there were

at least three individuals with SMI values. We excluded populations that had fewer than three site-surveys with mean SMI values. After processing, this dataset contained 1837 site-survey records.

2.2.1 | Site-Survey Analysis

To investigate the relationship between body condition and relative abundance, we fitted linear mixed models using the site-survey dataset (Figure 1). We used the log(mean SMI) of each site-survey as the response variable and inversely weighted the model with the SMI standard error. We used site-survey reptile capture rates as the predictor variable in models. Capture rates were scaled at the species level per dataset to standardize capture rates across studies (mean of zero and standard deviation of one). We included “dataset ID,” “site” nested within “dataset ID,” “species” nested within “dataset ID” and a phylogenetic distance matrix as random effects in the model to account for potential non-independence within and across the underlying datasets. We created the phylogenetic matrix using all study species from phylogeny trees obtained from the Vertlife database (Tonini et al. 2016). We first fitted a model with the site level capture rate as the sole predictor variable.

We fitted a second model investigating the interaction between capture rates and “disturbance” (levels: yes, no) to compare the relationship between relative abundance and body condition in disturbed sites and reference sites. We used the same random effects structure as in the previous model and a subset of the data that included populations with three or more site-surveys within a “disturbance” level. We fitted site-survey models using the Almer package (Bolstad et al. 2014). These models are unable to account for imperfect detection within the underlying datasets. Data and R code are available in Dryad (Macdonald 2025).

2.3 | Population Response Dataset

To investigate the relationship between body condition and relative abundance responses to disturbance, we compiled a population response dataset (Figure 1). For each species within each




FIGURE 1 | Overview of how the site-survey and population response datasets were created from original studies. LRR: Log-Response Ratios, SMI: Scaled Mass Index.

study, we calculated log-response ratios (LRR) for both SMI and capture rates to standardize our response and predictor variables across the underlying datasets. We calculated LRRs and their associated variances using the SingleCaseES package v0.7.2 (Pustejovsky et al. 2023). LRRs reflect the proportional change in response metrics between disturbance and reference values. For example, a negative abundance LRR reflects lower mean relative abundance within disturbed sites compared to reference sites, while a positive abundance LRR reflects greater mean relative abundance within disturbed sites compared to reference sites.

We excluded populations that had less than five reference or disturbed individuals. When datasets contained distinct repeat sampling (e.g., yearly surveys), LRRs were calculated across sites and individuals for each survey period independently. However, in one study (SW_01) there was one reference and one disturbance site, surveyed for three years. As such, we treated data from years as replicates, like sites in other datasets, and calculated LRRs by comparing the three unburnt surveys with the three burnt surveys.

When individual datasets included secondary variables which could introduce potential variation in the responses (e.g., vegetation type), we controlled for those secondary variables using the *aggregate* function within the LRR calculation (Pustejovsky et al. 2023). Likewise, when datasets contained a crossed design with multiple disturbance types, we calculated LRRs for each disturbance type with the other disturbance types controlled for within the *aggregate* function. After processing, this dataset contained 282 pairs of LRRs, created from 42,703 records.

2.3.1 | Population Response Analyses

We fitted meta-regression models to investigate the relationship between abundance responses to disturbance, and body condition responses to disturbance. We used a robust variance estimator, contained the degrees of freedom to the number of datasets (33) and used restricted maximum likelihood. To account for potential non-independence across effect sizes, we included the random effects; “dataset ID,” “species,” “species” nested within “dataset ID” and a phylogenetic distance matrix (Tonini et al. 2016) in each model. The capture rate LRR and associated variances were used as the response variable in our models. We included the linear and quadratic terms of body condition LRR as the predictor variables to investigate the relationship between these response metrics.

We subsequently fitted models that investigated how capture rate LRRs (response variable) were affected by the interaction between the linear or quadratic body condition LRR term and additional predictor variables. Additional predictor variables were disturbance group (levels: pulse, press), time since disturbance (pulse disturbance only), reproduction mode (levels: live bearing, egg laying), species body size (snout-vent length) and disturbance type (Table 1). For disturbance type, we limited the dataset to disturbances that contained more than 15 effect sizes from more than three datasets (combined), namely fire, habitat fragmentation, invasive species, and mining restoration. Most

of our studies investigated the impacts of fire (44% LRRs, 43% studies). Therefore, we ran an additional model investigating the interaction between body condition LRR and time since disturbance for fire data only.

To evaluate whether abundance responses to disturbance were related to body condition responses to disturbance from the previous year, we subsetted the population response dataset to include only studies with repeat annual surveys. We then paired population abundance LRR with the body condition LRR from the previous year. This dataset contained 69 pairs of LRRs, created from 10,250 records across 10 studies. Using the same model structure, we fitted meta-regression models using the body condition LRR from the previous year as the predictor variable. We used the *metafor* package v3.8-1 (Viechtbauer 2010) to fit all meta-regression models. These models are unable to account for imperfect detection within the underlying datasets. Data and R code are available in Dryad (Macdonald 2025).

2.3.2 | Sensitivity Analysis

To assess the robustness of our population response analysis, we ran two types of sensitivity analyses on the meta-regression models: Cook’s distance outliers and body condition variance. When results were significant ($p < 0.05$) we assessed the influence of outliers using Cook’s distance values by refitting the models without each effect size that had a Cook’s distance value > 1 and again without all effect sizes with a Cook’s distance > 1 (Viechtbauer and Cheung 2010).

In the population response dataset, each SMI LRR value had an associated variance that could not be accounted for in the meta-regression models. For body condition variance sensitivity, we refitted the models after excluding data with large SMI LRR variance (mean + 2 SD), that is, lower confidence.

3 | Results

3.1 | Site-Survey Responses

We found a significant positive relationship between the mean Scaled Mass Index (SMI) and scaled capture rates at the site-survey level ($\beta \pm SE = 0.009 \pm 0.003$, $t = 3.22$, $p = 0.002$), whereby an increase in capture rates of one standard deviation corresponds with a SMI increase of one gram. We found that mean SMI was significantly lower in disturbed sites compared to undisturbed sites ($F_{1,39} = 8.45$, $p = 0.006$). However, we found no evidence for an interactive effect between scaled capture rates and disturbance (levels: yes, no) ($F_{1,39} = 0.406$, $p = 0.528$).

3.2 | Population Responses

At the population level, we found a relatively even spread of congruent responses (where disturbance had either a positive or negative effect on both mean abundance and mean body condition) and contrasting responses (where disturbance had an opposing effect on the two metrics) (Figure 2a).

TABLE 1 | Disturbance characteristics and species traits, with justifications on how they may impact abundance and body condition responses to disturbance.

Variable	Justification
Disturbance group levels: press, pulse. Press: Fragmentation, Grazing, Invasive Species, Land cover change, Urbanisation, Wind turbines. Pulse: Drought/Low rainfall, Fire, Logging, Logging/Fire, Mining rehabilitation	Lizards can adapt to disturbed landscapes and regain lost body condition (Rodríguez-Prieto et al. 2010). Therefore, we might expect weaker body condition responses following disturbances which are ongoing (i.e., press disturbances) compared to pulse disturbances which are acute, and which habitats can recover from
Time since disturbance (years). Pulse disturbances only	Resource availability can change with time since disturbance, often defined by post disturbance successional trajectories (Hutchinson 1941)
Disturbance type levels: Drought/Low rainfall, Edge, Fire, Fragmentation, Grazing, Invasive Species, Land cover change, Logging, Logging/Fire, Mining rehabilitation, Urbanisation, Wind turbines	Reptile abundance and body condition responses can differ across disturbance types (Doherty et al. 2020; Macdonald et al. 2023). The influence of disturbance may differ between metrics. For example, disturbance types that destroy food resources may have greater influence on body condition than abundance
Reproduction mode (levels: live bearing, egg laying)	Macdonald et al. (2024; unpublished data) found similarities in the body condition and abundance responses of two live bearing species. Live bearing species have greater basking and foraging requirements, lower reproduction rates, and slower population recovery compared to egg laying species (Lazzari et al. 2022; Meiri et al. 2012; Robert and Thompson 2007). Therefore, disturbances may have larger effects on the body condition of live bearing species compared to egg laying species
Body size (mean species snout-vent length)	Although it was not an important predictor of reptile body condition or abundance responses to disturbance at the population level (Doherty et al. 2020; Macdonald et al. 2023), it is not known whether body size has a deterministic effect on the relationship between abundance and body condition in response to disturbance. Smaller species may be more likely to survive disturbance events (Böhm et al. 2016), but the reduction in resources and limited ability to emigrate may increase intra-specific competition for small species, while any surviving individuals of large species may have increased mobility and access to potential resources




FIGURE 2 | (a) Log-response ratios of abundance and body condition responses to disturbance and 85% confidence intervals. Disturbance had a negative (red), positive (green) or variable (blue) effects on abundance and body condition. (b) Body condition variance sensitivity analysis investigating the quadratic body condition log-response ratios (LRR) against abundance LRR. Dataset excludes effect sizes with large body condition LRR variance (mean + 2 SD). Dashed lines: 95% confidence intervals.

We found no overall relationship between abundance log-response ratios (LRR) and linear ($\beta \pm \text{SE} = 0.323 \pm 0.47$, $p = 0.495$) or quadratic ($\beta \pm \text{SE} = 1.620 \pm 3.48$, $p = 0.645$) body condition LRR. However, when assessing body condition

variance sensitivity by excluding results with high SD (see Section 2.3.2 Sensitivity Analysis), we found a significant non-linear relationship between SMI LRRs and abundance LRRs, whereby larger body condition responses to disturbance

(both positive and negative) were associated with negative abundance responses to disturbance ($\beta \pm \text{SE} = -4.324 \pm 1.90$, $p = 0.030$, Figure 2b).

3.2.1 | Disturbance Group

We found no evidence for a linear relationship between abundance and body condition LRRs when disturbance group (press/pulse) was included in the model ($\beta \pm \text{SE} = -0.002 \pm 1.19$, $p = 0.999$), nor when assessing body condition variance sensitivity (see Section 2.3.2 Sensitivity Analysis).

Abundance LRR was significantly affected by a non-linear interactive effect between body condition LRR and disturbance group, whereby larger body condition responses to disturbances were associated with positive abundance responses to press disturbances, but with negative abundance responses to pulse disturbances ($\beta \pm \text{SE} = -15.91 \pm 2.62$, $p < 0.001$, Figure S1). When excluding Cook's distance outliers (see Section 2.3.2 Sensitivity Analysis), the inference of the results remained the same. However, the relationship was lost when assessing body condition variance sensitivity (see Section 2.3.2 Sensitivity Analysis).

3.2.2 | Disturbance Type

We found no linear relationship between abundance and body condition LRRs when we included disturbance type in the model ($F_{7,19} = 1.22$, $p = 0.339$), including when we assessed sensitivity to body condition variance (see Section 2.3.2 Sensitivity Analysis).

We found a significant non-linear interactive effect between body condition LRR and disturbance type, whereby larger positive and negative body condition LRRs for fragmentation and invasive species were associated with positive abundance LRRs ($F_{6,15} = 15.28$, $p < 0.001$, Figure S2). The inference of the results remained largely the same when individual Cook's distance outliers were removed (see Section 2.3.2 Sensitivity Analysis). However, the relationship disappeared when all Cook's distance outliers were removed and when assessing body condition variance sensitivity (see Section 2.3.2 Sensitivity Analysis).

3.2.3 | Time Since Disturbance

We found no linear or quadratic relationship of abundance LRR with the interaction between body condition LRR and time since disturbance, nor in any sensitivity analyses (linear: $\beta \pm \text{SE} = 0.06 \pm 0.21$, $p = 0.778$, quadratic: $\beta \pm \text{SE} = 1.17 \pm 1.37$, $p = 0.421$). Likewise, we found no relationships between abundance and body condition LRRs with time since disturbance for fire datasets only (linear: $\beta \pm \text{SE} = 0.02 \pm 0.21$, $p = 0.923$, quadratic: $\beta \pm \text{SE} = 1.07 \pm 1.22$, $p = 0.411$).

3.2.4 | Species Traits

We found no relationships between abundance and body condition LRRs when investigating the interactive effects with body size (linear: $\beta \pm \text{SE} = 0.003 \pm 0.003$, $p = 0.402$, quadratic:

$\beta \pm \text{SE} = -0.02 \pm 0.02$, $p = 0.306$), or reproductive mode (linear: $\beta \pm \text{SE} = 0.63 \pm 1.27$, $p = 0.623$, quadratic: $\beta \pm \text{SE} = -0.25 \pm 3.18$, $p = 0.937$), nor in any sensitivity analyses.

3.2.5 | Previous Years' Body Condition

We found no evidence for a relationship between abundance LRR and body condition LRR from the previous year (linear: $\beta \pm \text{SE} = 0.62 \pm 0.727$, $p = 0.421$, quadratic: $\beta \pm \text{SE} = -8.94 \pm 7.69$, $p = 0.283$), nor in any sensitivity analyses.

4 | Discussion

Our results reveal a positive relationship between reptile capture rates and body condition at the site-survey level. The implication of these results is that population size is more likely to be driven by individual fitness compared to density dependence (Herrando-Pérez et al. 2012). These results align with some past research that found lower reptile body condition was associated with lower reproductive output and higher mortality risk (Reading 2004; Shine et al. 2001). However, the results were also highly variable, which aligns with research by Lee-Yaw et al. (2022) who found that species distribution models had poor predictive capacity for a range of community, population, and individual response metrics. Overall, our site-survey results provide only limited support for our first prediction. They also provide limited support for the population collapse framework by showing a concurrent relationship between the two metrics (Cerini et al. 2023).

Environmental variables may weaken the relationship between body condition and abundance by having strong but variable influences on each response metric. For example, in Australia, *Gehyra variegata* (gecko) population size and body condition were strongly influenced by different climatic factors; summer temperature and rainfall were important drivers of population size, whereas winter rainfall and river height were important drivers of body condition variation (Grimm-Seyfarth et al. 2018). Such results highlight the potential for idiosyncratic responses across studies to weaken the expected positive relationship between body condition and abundance.

4.1 | Population Responses to Disturbance

We found little evidence for a relationship between reptile abundance and body condition responses to disturbance (Figure 2). Our population results provided no support for our second or third predictions and, as such, no systematic overall support for the risk-disturbance hypothesis (Frid and Dill 2002) or the population collapse framework (Cerini et al. 2023). Disturbances may modify or uncouple the relationship between abundance and body condition by influencing underlying drivers, including predation, competition, and resource availability (Figure 3). These drivers can, in turn, influence species behavior (e.g., foraging and dispersal) and mortality risk (Figure 3). For example, fire can alter predator-prey dynamics (Doherty et al. 2022) which may, in turn, influence population size via direct predation and influence




FIGURE 3 | Hypothesized causal links between reptile body condition and abundance and the additional influence of resources, competitors, and predators on this relationship. Disturbances and other abiotic factors can also influence these pathways and interactions. Solid arrows indicate direct links, while dashed arrows indicate indirect links. ¹Heithaus et al. (2007), ²Amo et al. (2006), ³Stewart et al. (2005), ⁴Keyser et al. (2005), ⁵Catling (1988), ⁶Lannoo and Stiles (2017), ⁷Reading (2004), ⁸Shine et al. (2001), ⁹Walkup et al. (2017).

body condition via changes in foraging behavior or competition (Figure 3). Likewise, recent work found that invasive predator management could have a strong influence on reptile abundance (unpublished data from Macdonald et al.), while having little effect on reptile body condition (Macdonald et al. 2024). These contrasting results could be explained by greater predator densities influencing the behavior and densities of target species and their competitors, while changes in direct predation concurrently reduced intra-specific competition (Figure 3). Future research should examine the key drivers influencing the relationships between response metrics, particularly predators, competitors, and changes in resource availability (Figure 3). Understanding how key drivers influence population carrying capacity and via which mechanisms (e.g., changes in population size, foraging, thermoregulation

or dispersal) may improve predictive frameworks that are important in conservation management (Cross et al. 2021; Gibbons et al. 2000; Pulsford et al. 2016). This work would be especially valuable for species where disturbance is known to lead to population collapse (Walkup et al. 2017). These mechanisms could be explored using natural field experiments as well as experimental approaches (e.g., in mesocosms or enclosures) that control resource availability or population densities under disturbed conditions (Baruah et al. 2019).

Disturbances can alter the relationship between response metrics. For example, Sinclair et al. (2023) found that anthropogenic effects (commercial harvest, nutrient availability, water transparency, and temperature) reversed the relationship between response metrics (size-at-age and population size) by

altering intra-specific competition pressure. Likewise, Rossi and Leiner (2022) revealed that density feedback relationships between small mammal population size and growth were affected by fire, intra-specific competition, and additional environmental variables. Additional environmental factors may have a compounding or competing influence on the relationship between response metrics and the underlying mechanisms, interactions, and feedback loops. As such, predicting outcomes of a disturbance on the relationship between body condition and population size is complex.

We found some evidence that disturbance characteristics can influence the relationship between body condition and abundance responses to disturbance. However, the relationships were not robust to sensitivity analyses, highlighting the uncertainty of these links. If they are real effects, the implication is that the underlying mechanisms of disturbances can change intra-specific competition pressure, thereby altering the relationship between abundance and body condition. For example, pulse disturbances (e.g., fire, logging, drought) can greatly reduce resources for some species, which could increase intra-specific competition resulting in both lower body condition and abundance. Pulse disturbances can also result in large direct mortality for some species, which would lower abundance, but body condition may increase due to reduced intra-specific competition between surviving individuals. Under press disturbances, some species are more competitive (e.g., edge specialists, habitat generalists) which could increase carrying capacity, resulting in both higher abundance and body condition. In contrast, under press disturbances such as fragmentation, the abundance of some species may increase within the remaining fragments, thereby increasing intraspecific competition leading to poorer body condition (Ostfeld 1994). It will be important to continue investigating how disturbance characteristics influence important underlying mechanisms that may drive changes in different response metrics and their relationship.

In addition, future research should investigate whether combined abundance and body condition responses can provide better insight into disturbance impacts and extinction risk. For example, if a population has lower abundance and the persisting individuals have low body condition, is that population at a greater risk of ongoing declines compared to a population with low abundance but better body condition? Alternatively, researchers may ask, is poor body condition a symptom of severe population decline and, therefore, too difficult to correlate with population decline until it is too late for conservation intervention?

5 | Conclusion

Our results suggest that changes in body condition in response to disturbance cannot generally be used to predict how species abundance responds to disturbance. As such, we found little support for the risk disturbance hypothesis or population collapse framework. Future research and monitoring programs investigating the impacts of disturbance on body condition should be careful not to infer population-level responses without the relationship being confirmed for their study population. We highlight important ecological pathways between disturbance, predators, competitors, and resources, and their potential to

disrupt the relationship between abundance and body condition responses (Figure 3). We encourage further investigation into these key drivers to empirically test and advance ecological theory.

Author Contributions

Kristina J. Macdonald: conceptualization, data curation, formal analysis, investigation, methodology, project administration, visualization, writing – original draft, writing – review and editing. **Don A. Driscoll:** conceptualization, funding acquisition, methodology, supervision, writing – review and editing. **Michael D. Craig:** investigation, writing – review and editing. **Robert A. Davis:** investigation, writing – review and editing. **Steven J. Hromada:** investigation, writing – review and editing. **C. M. Gienger:** investigation, writing – review and editing. **Lee A. Fitzgerald:** investigation, writing – review and editing. **Daniel J. Leavitt:** investigation, writing – review and editing. **Danielle K. Walkup:** data curation, investigation, writing – review and editing. **Rickard Abom:** investigation, writing – review and editing. **Adrian F. Wayne:** investigation, writing – review and editing. **Gordon R. Friend:** investigation, writing – review and editing. **Brent Johnson:** investigation, writing – review and editing. **Danielle Stokeld:** investigation, writing – review and editing. **Chris R. Dickman:** investigation, writing – review and editing. **Scott A. Thompson:** investigation, writing – review and editing. **Graham G. Thompson:** investigation, writing – review and editing. **Daniel F. Bohórquez Fandiño:** investigation, writing – review and editing. **Christopher K. Woolley:** investigation, writing – review and editing. **Annabel L. Smith:** investigation, writing – review and editing. **Calum Irvine:** investigation, writing – review and editing. **Tim S. Jessop:** investigation, writing – review and editing. **Jade E. Keehn:** investigation, writing – review and editing. **Chris R. Feldman:** investigation, writing – review and editing. **Xavier Santos:** investigation, writing – review and editing. **Josabel Belliure:** investigation, writing – review and editing. **Juli G. Pausas:** investigation, writing – review and editing. **Stacey L. Weiss:** investigation, writing – review and editing. **Patricia A. Fleming:** investigation, writing – review and editing. **Shannon Dundas:** investigation, writing – review and editing. **Dylan M. Westaway:** investigation, writing – review and editing. **Sabrina E. Duncan:** investigation, writing – review and editing. **Tim S. Doherty:** conceptualization, investigation, methodology, supervision, writing – review and editing.

Acknowledgements

We acknowledge the various funders who contributed to the underlying datasets used in this research including the Australian Research Council (LP170101134, AD9031596), the Australian Research Grant Scheme (A18615230), Barrick Kanowna, Charles Darwin University, Edith Cowan University, Holsworth Wildlife Research Endowment, Minerals Research Institute of Western Australia, M.J. Murdock Charitable Trust, Australian Government's National Environmental Science Program, Northern Australian Environmental Resources Hub, Norilsk Nickel, Northern Territory Department of Lands, Planning and Environment, OMG Cawse Nickel Operations, Parks Victoria, Placer Dome Asia Pacific Kalgoorlie West Operations, and the Threatened Species Recovery Hub. C.R.D. thanks T and L Churches for access to Mooraberrie. We thank the anonymous reviewers for their feedback on an earlier version of this manuscript. Open access publishing facilitated by Charles Darwin University, as part of the Wiley & hyphen; Charles Darwin University agreement via the Council of Australian University Librarians.

Conflicts of Interest

The authors declare no conflicts of interest.

Data Availability Statement

The data that support the findings of this study are openly available in Dryad at <https://doi.org/10.5061/dryad.wdbrv160k>.

References

- Amo, L., P. López, and J. Martín. 2006. "Nature-Based Tourism as a Form of Predation Risk Affects Body Condition and Health State of *Podarcis muralis* Lizards." *Biological Conservation* 131, no. 3: 402–409. <https://doi.org/10.1016/j.biocon.2006.02.015>.
- Aubry, L. M., R. F. Rockwell, E. G. Cooch, R. W. Brook, C. P. H. Mulder, and D. N. Koons. 2013. "Climate Change, Phenology, and Habitat Degradation: Drivers of Gosling Body Condition and Juvenile Survival in Lesser Snow Geese." *Global Change Biology* 19, no. 1: 149–160. <https://doi.org/10.1111/gcb.12013>.
- Baruah, G., C. F. Clements, F. Guillaume, and A. Ozgul. 2019. "When Do Shifts in Trait Dynamics Precede Population Declines?" *American Naturalist* 193, no. 5: 633–644. <https://doi.org/10.1086/702849>.
- Battisti, C., G. Poeta, and G. Fanelli. 2016. "The Concept of Disturbance." In *An Introduction to Disturbance Ecology: A Road Map for Wildlife Management and Conservation*, edited by C. Battisti, G. Poeta, and G. Fanelli, 7–12. Springer International Publishing. https://doi.org/10.1007/978-3-319-32476-0_2.
- Böhm, M., R. Williams, H. R. Bramhall, et al. 2016. "Correlates of Extinction Risk in Squamate Reptiles: The Relative Importance of Biology, Geography, Threat and Range Size." *Global Ecology and Biogeography* 25, no. 4: 391–405. <https://doi.org/10.1111/geb.12419>.
- Bolstad, G. H., T. F. Hansen, C. Pélabon, M. Falahati-Anbaran, R. Pérez-Barrales, and W. S. Armbruster. 2014. "Genetic Constraints Predict Evolutionary Divergence in *Dalechampia* Blossoms." *Philosophical Transactions of the Royal Society, B: Biological Sciences* 369, no. 1649: 20130255. <https://doi.org/10.1098/rstb.2013.0255>.
- Catling, P. C. 1988. "Similarities and Contrasts in the Diets of Foxes, Vulpes Vulpes, and Cats, *Felis catus*, Relative to Fluctuating Prey Populations and Drought." *Wildlife Research* 15, no. 3: 307. <https://doi.org/10.1071/WR9880307>.
- Cerini, F., D. Z. Childs, and C. F. Clements. 2023. "A Predictive Timeline of Wildlife Population Collapse." *Nature Ecology & Evolution* 7, no. 3: 320–331. <https://doi.org/10.1038/s41559-023-01985-2>.
- Cogălniceanu, D., F. Stănescu, D. Székely, T. S. Topliceanu, R. Losif, and P. Székely. 2021. "Age, Size and Body Condition Do Not Equally Reflect Population Response to Habitat Change in the Common Spadefoot Toad *Pelobates fuscus*." *PeerJ* 9: e11678. <https://doi.org/10.7717/PEERJ.11678>.
- Cordier, J. M., R. Aguilar, J. N. Lescano, et al. 2021. "A Global Assessment of Amphibian and Reptile Responses to Land-Use Changes." *Biological Conservation* 253: 108863. <https://doi.org/10.1016/j.biocon.2020.108863>.
- Cox, N., B. E. Young, P. Bowles, et al. 2022. "A Global Reptile Assessment Highlights Shared Conservation Needs of Tetrapods." *Nature* 605, no. 7909: 285–290. <https://doi.org/10.1038/s41586-022-04664-7>.
- Cross, S. L., A. T. Cross, S. Tomlinson, S. M. Clark-Ioannou, P. G. Nevill, and P. W. Bateman. 2021. "Mitigation and Management Plans Should Consider All Anthropogenic Disturbances to Fauna." *Global Ecology and Conservation* 26: e01500. <https://doi.org/10.1016/j.gecco.2021.e01500>.
- Deng, S., C. M. Beale, P. J. Platts, and C. D. Thomas. 2023. "Human Modification of Land Cover Alters Net Primary Productivity, Species Richness and Their Relationship." *Global Ecology and Biogeography* 33: 385–399. <https://doi.org/10.1111/geb.13795>.
- Doherty, T. S., S. Balouch, K. Bell, et al. 2020. "Reptile Responses to Anthropogenic Habitat Modification: A Global Meta-Analysis." *Global Ecology and Biogeography* 29, no. 7: 1265–1279. <https://doi.org/10.1111/geb.13091>.
- Doherty, T. S., W. L. Geary, C. J. Jolly, et al. 2022. "Fire as a Driver and Mediator of Predator–Prey Interactions." *Biological Reviews* 97, no. 4: 1539–1558. <https://doi.org/10.1111/brv.12853>.
- Emery, J.-P., L. E. Valentine, Y. Hitchen, and N. Mitchell. 2021. "Survival of an Extinct in the Wild Skink From Christmas Island Is Reduced by an Invasive Centipede: Implications for Future Reintroductions." *Biological Invasions* 23, no. 2: 581–592. <https://doi.org/10.1007/s10530-020-02386-3>.
- Fair, W. S., and S. E. Henke. 1997. "Effects of Habitat Manipulations on Texas Horned Lizards and Their Prey." *Journal of Wildlife Management* 61, no. 4: 1366. <https://doi.org/10.2307/3802139>.
- Frid, A., and L. Dill. 2002. "Human-Caused Disturbance Stimuli as a Form of Predation Risk." *Conservation Ecology* 6, no. 1: 11. <https://doi.org/10.5751/es-00404-060111>.
- Gardner, J. L., T. Amano, W. J. Sutherland, M. Clayton, and A. Peters. 2016. "Individual and Demographic Consequences of Reduced Body Condition Following Repeated Exposure to High Temperatures." *Ecology* 97, no. 3: 786–795. <https://doi.org/10.1890/15-0642.1>.
- Gibbons, J. W., D. E. Scott, T. J. Ryan, et al. 2000. "The Global Decline of Reptiles, Déjà Vu Amphibians: Reptile Species Are Declining on a Global Scale. Six Significant Threats to Reptile Populations Are Habitat Loss and Degradation, Introduced Invasive Species, Environmental Pollution, Disease, Unsustainable Use, and Global Climate Change." *Bioscience* 50, no. 8: 653–666. [https://doi.org/10.1641/0006-3568\(2000\)050\[0653:TGDORD\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2000)050[0653:TGDORD]2.0.CO;2).
- Gould, P. R., K. K. Cecala, S. S. Drukker, B. A. McKenzie, and C. H. Van de Ven. 2024. "Should Landscape Variation in Population Status Be Assessed With Individual- Or Population-Level Indicators?" *Journal of Wildlife Management* 88, no. 2: e22520. <https://doi.org/10.1002/jwmg.22520>.
- Grimm-Seyfarth, A., J.-B. Mihoub, B. Gruber, and K. Henle. 2018. "Some Like It Hot: From Individual to Population Responses of an Arboreal Arid-Zone Gecko to Local and Distant Climate." *Ecological Monographs* 88, no. 3: 336–352. <https://doi.org/10.1002/ecm.1301>.
- Heithaus, M. R., A. Frid, A. J. Wirsing, et al. 2007. "State-Dependent Risk-Taking by Green Sea Turtles Mediates Top-Down Effects of Tiger Shark Intimidation in a Marine Ecosystem." *Journal of Animal Ecology* 76, no. 5: 837–844. <https://doi.org/10.1111/j.1365-2656.2007.01260.x>.
- Herrando-Pérez, S., S. Delean, B. W. Brook, and C. J. A. Bradshaw. 2012. "Density Dependence: An Ecological Tower of Babel." *Oecologia* 170, no. 3: 585–603. <https://doi.org/10.1007/s00442-012-2347-3>.
- Hillebrand, H., and C. Kunze. 2020. "Meta-Analysis on Pulse Disturbances Reveals Differences in Functional and Compositional Recovery Across Ecosystems." *Ecology Letters* 23, no. 3: 575–585. <https://doi.org/10.1111/ele.13457>.
- Hutchinson, G. E. 1941. "Ecological Aspects of Succession in Natural Populations." *American Naturalist* 75, no. 760: 406–418. <https://doi.org/10.1086/280983>.
- IPBES. 2019. "Global Assessment Report on Biodiversity and Ecosystem Services of the Intergovernmental Science—Policy Platform on Biodiversity and Ecosystem Services." In *Zenodo*, edited by E. S. Brondizio, J. Settele, S. Díaz, and H. T. Ngo, 1–1082. IPBES secretariat. <https://doi.org/10.5281/ZENODO.6417333>.
- Janin, A., J.-P. Léna, and P. Joly. 2011. "Beyond Occurrence: Body Condition and Stress Hormone as Integrative Indicators of Habitat Availability and Fragmentation in the Common Toad." *Biological Conservation* 144, no. 3: 1008–1016. <https://doi.org/10.1016/j.biocon.2010.12.009>.
- Keehn, J. E., K. T. Shoemaker, and C. R. Feldman. 2019. "Population-Level Effects of Wind Farms on a Desert Lizard." *Journal of Wildlife Management* 83, no. 1: 145–157. <https://doi.org/10.1002/jwmg.21565>.
- Keinath, D. A., D. F. Doak, K. E. Hodges, et al. 2017. "A Global Analysis of Traits Predicting Species Sensitivity to Habitat Fragmentation." *Global Ecology and Biogeography* 26, no. 1: 115–127. <https://doi.org/10.1111/geb.12509>.
- Keyser, P. D., D. C. Guynn, and H. S. Hill. 2005. "Population Density–Physical Condition Relationships in White-Tailed Deer." *Journal of*

- Wildlife Management* 69, no. 1: 356–365. [https://doi.org/10.2193/0022-541X\(2005\)069<0356:PDCRIW>2.0.CO;2](https://doi.org/10.2193/0022-541X(2005)069<0356:PDCRIW>2.0.CO;2).
- Labocha, M. K., H. Schutz, and J. P. Hayes. 2014. “Which Body Condition Index Is Best?” *Oikos* 123, no. 1: 111–119. <https://doi.org/10.1111/j.1600-0706.2013.00755.x>.
- Laidre, K. L., J. A. Estes, M. T. Tinker, J. Bodkin, D. Monson, and K. Schneider. 2006. “Patterns of Growth and Body Condition in Sea Otters From the Aleutian Archipelago Before and After the Recent Population Decline.” *Journal of Animal Ecology* 75, no. 4: 978–989. <https://doi.org/10.1111/j.1365-2656.2006.01117.x>.
- Lannoo, M. J., and R. M. Stiles. 2017. “Effects of Short-Term Climate Variation on a Long-Lived Frog.” *Copeia* 105, no. 4: 726–733. <https://doi.org/10.1643/CH-16-449>.
- Lazzari, J., C. F. Sato, and D. A. Driscoll. 2022. “Traits Influence Reptile Responses to Fire in a Fragmented Agricultural Landscape.” *Landscape Ecology* 37, no. 9: 2363–2382. <https://doi.org/10.1007/s10980-022-01417-9>.
- Lee-Yaw, J. A., J. L. McCune, S. Pironon, and S. N. Sheth. 2022. “Species Distribution Models Rarely Predict the Biology of Real Populations.” *Ecography* 2022: e05877. <https://doi.org/10.1111/ecog.05877>.
- Legge, S., H. Bijlani, B. Taylor, et al. 2024. “Pirra Jungku and Pirra Warlu: Using Traditional Fire-Practice Knowledge and Contemporary Science to Guide Fire-Management Goals for Desert Animals.” *Wildlife Research* 51, no. 10: WR24069. <https://doi.org/10.1071/WR24069>.
- Lunghi, E., R. Manenti, M. Mulargia, M. Veith, C. Corti, and G. F. Ficetola. 2018. “Environmental Suitability Models Predict Population Density, Performance and Body Condition for Microendemic Salamanders.” *Scientific Reports* 8, no. 1: 7527. <https://doi.org/10.1038/s41598-018-25704-1>.
- Macdonald, K. J. 2025. “Data and Code: Cross-Continental Analysis Shows That Disturbance Effects on Reptile Body Condition Do Not Predict Abundance Responses.” *Dryad*. <https://doi.org/10.5061/dryad.wdbrv160k>.
- Macdonald, K. J., T. S. Doherty, B. A. Hradsky, and D. A. Driscoll. 2024. “Disturbance and Ecosystem Management Interact to Shape Reptile Body Condition.” *Animal Conservation* 28: 12983. <https://doi.org/10.1111/acv.12983>.
- Macdonald, K. J., D. A. Driscoll, K. J. Macdonald, B. Hradsky, and T. S. Doherty. 2023. “Meta-Analysis Reveals Impacts of Disturbance on Reptile and Amphibian Body Condition.” *Global Change Biology* 29, no. 17: 4949–4965. <https://doi.org/10.1111/gcb.16852>.
- Madsen, T., J. Loman, D. Bauwens, et al. 2023. “The Impact of an Extreme Climatic Event on Adder (*Vipera berus*) Demography in Southern Sweden.” *Biological Journal of the Linnean Society* 138: blac147. <https://doi.org/10.1093/biolinnean/blac147>.
- Meiri, S., J. H. Brown, and R. M. Sibly. 2012. “The Ecology of Lizard Reproductive Output.” *Global Ecology and Biogeography* 21, no. 5: 592–602. <https://doi.org/10.1111/j.1466-8238.2011.00700.x>.
- Ostfeld, R. S. 1994. “The Fence Effect Reconsidered.” *Oikos* 70, no. 3: 340–348. <https://doi.org/10.2307/3545771>.
- Parmesan, C., M. Morecroft, Y. Trisurat, et al. 2022. *IPCC AR6 WGII Chapter 2—Terrestrial and Freshwater Ecosystems and Their Services*, 197–377. Cambridge University Press. <https://doi.org/10.1017/9781009325844.004>.
- Pausas, J. G., and J. E. Keeley. 2009. “A Burning Story: The Role of Fire in the History of Life.” *Bioscience* 59, no. 7: 593–601. <https://doi.org/10.1525/bio.2009.59.7.10>.
- Peig, J., and A. J. Green. 2009. “New Perspectives for Estimating Body Condition From Mass/Length Data: The Scaled Mass Index as an Alternative Method.” *Oikos* 118, no. 12: 1883–1891. <https://doi.org/10.1111/j.1600-0706.2009.17643.x>.
- Peig, J., and A. J. Green. 2010. “The Paradigm of Body Condition: A Critical Reappraisal of Current Methods Based on Mass and Length.” *Functional Ecology* 24, no. 6: 1323–1332. <https://doi.org/10.1111/j.1365-2435.2010.01751.x>.
- Pigeon, G., T. H. G. Ezard, M. Festa-Bianchet, D. W. Coltman, and F. Pelletier. 2017. “Fluctuating Effects of Genetic and Plastic Changes in Body Mass on Population Dynamics in a Large Herbivore.” *Ecology* 98, no. 9: 2456–2467. <https://doi.org/10.1002/ecy.1940>.
- Pulsford, S. A., D. B. Lindenmayer, and D. A. Driscoll. 2016. “A Succession of Theories: Purging Redundancy From Disturbance Theory.” *Biological Reviews* 91, no. 1: 148–167. <https://doi.org/10.1111/brv.12163>.
- Pustejovsky, J. E., M. Chen, and D. M. Swan. 2023. “SingleCaseES: A Calculator for Single-Case Effect Sizes (Version 0.7.2) [R Package].” <https://jepusto.github.io/SingleCaseES/>.
- Quesnelle, P. E., K. E. Lindsay, and L. Fahrig. 2014. “Low Reproductive Rate Predicts Species Sensitivity to Habitat Loss: A Meta-Analysis of Wetland Vertebrates.” *PLoS One* 9, no. 3: e90926. <https://doi.org/10.1371/JOURNAL.PONE.0090926>.
- R Core Team. 2022. *R: A Language and Environment for Statistical Computing (4.2.1)*. R Foundation for Statistical Computing. <https://www.R-project.org/>.
- Reading, C. J. 2004. “The Influence of Body Condition and Prey Availability on Female Breeding Success in the Smooth Snake (*Coronella austriaca* Laurenti).” *Journal of Zoology* 264, no. 1: 61–67. <https://doi.org/10.1017/S0952836904005515>.
- Reading, C. J. 2007. “Linking Global Warming to Amphibian Declines Through Its Effects on Female Body Condition and Survivorship.” *Oecologia* 151, no. 1: 125–131. <https://doi.org/10.1007/s00442-006-0558-1>.
- Robert, K. A., and M. B. Thompson. 2007. “Is Basking Opportunity in the Viviparous Lizard, *Eulamprus tympanum*, Compromised by the Presence of a Predator Scent?” *Journal of Herpetology* 41, no. 2: 287–293. [https://doi.org/10.1670/0022-1511\(2007\)41\[287:IBOITV\]2.0.CO;2](https://doi.org/10.1670/0022-1511(2007)41[287:IBOITV]2.0.CO;2).
- Rodríguez-Prieto, I., J. Martín, and E. Fernández-Juricic. 2010. “Habituation to Low-Risk Predators Improves Body Condition in Lizards.” *Behavioral Ecology and Sociobiology* 64, no. 12: 1937–1945. <https://doi.org/10.1007/s00265-010-1004-2>.
- Rossi, R. C., and N. O. Leiner. 2022. “Weather, Fire, and Density Drive Population Dynamics of Small Mammals in the Brazilian Cerrado.” *Journal of Mammalogy* 103, no. 5: 1127–1140. <https://doi.org/10.1093/jmammal/gyac053>.
- Santos, J. L., H. Sitters, D. A. Keith, W. L. Geary, R. Tingley, and L. T. Kelly. 2022. “A Demographic Framework for Understanding Fire-Driven Reptile Declines in the ‘Land of the Lizards.’” *Global Ecology and Biogeography* 31, no. 10: 2105–2119. <https://doi.org/10.1111/geb.13520>.
- Schwarzkopf, L. 2022. “Homage to Reptiles and Amphibians as Model Systems: One Ecologist’s View.” *Journal of Herpetology* 56, no. 1: 45–55. <https://doi.org/10.1670/21-020>.
- Sergio, F., J. Blas, and F. Hiraldo. 2018. “Animal Responses to Natural Disturbance and Climate Extremes: A Review.” *Global and Planetary Change* 161: 28–40. <https://doi.org/10.1016/j.gloplacha.2017.10.009>.
- Shine, R., M. P. LeMaster, I. T. Moore, M. M. Olsson, and R. T. Mason. 2001. “Bumpus in the Snake Den: Effects of Sex, Size and Body Condition on Mortality of Red-Sided Garter Snakes.” *Evolution* 55, no. 3: 598–604. <https://doi.org/10.1111/j.0014-3820.2001.tb00792.x>.
- Sinclair, J. S., R. Briland, M. E. Fraker, et al. 2023. “Anthropogenic Change Decouples a Freshwater Predator’s Density Feedback.” *Scientific Reports* 13, no. 1: 7613. <https://doi.org/10.1038/s41598-023-34408-0>.
- Smith, A. L. 2018. “Successional Changes in Trophic Interactions Support a Mechanistic Model of Post-Fire Population Dynamics.” *Oecologia* 186, no. 1: 129–139. <https://doi.org/10.1007/s00442-017-4016-z>.

- Sousa, W. P. 1984. "The Role of Disturbance in Natural Communities." *Annual Review of Ecology and Systematics* 15: 353–391. <http://www.jstor.org/stable/2096953>.
- Stewart, J. D., J. W. Durban, H. Fearnbach, et al. 2021. "Survival of the Fattest: Linking Body Condition to Prey Availability and Survivorship of Killer Whales." *Ecosphere* 12, no. 8: e03660. <https://doi.org/10.1002/ecs2.3660>.
- Stewart, K. M., R. T. Bowyer, B. L. Dick, B. K. Johnson, and J. G. Kie. 2005. "Density-Dependent Effects on Physical Condition and Reproduction in North American Elk: An Experimental Test." *Oecologia* 143, no. 1: 85–93. <https://doi.org/10.1007/s00442-004-1785-y>.
- Testa, J. W., and G. P. Adams. 1998. "Body Condition and Adjustments to Reproductive Effort in Female Moose (*Alces alces*)." *Journal of Mammalogy* 79, no. 4: 1345–1354. <https://doi.org/10.2307/1383026>.
- Tingley, R., S. L. Macdonald, N. J. Mitchell, et al. 2019. "Geographic and Taxonomic Patterns of Extinction Risk in Australian Squamates." *Biological Conservation* 238: 108203. <https://doi.org/10.1016/j.biocon.2019.108203>.
- Tonini, J. F. R., K. H. Beard, R. B. Ferreira, W. Jetz, and R. A. Pyron. 2016. "Fully-Sampled Phylogenies of Squamates Reveal Evolutionary Patterns in Threat Status." *Biological Conservation* 204: 23–31. <https://doi.org/10.1016/j.biocon.2016.03.039>.
- Viechtbauer, W. 2010. "Conducting Meta-Analyses in R With the Meta for Package." *Journal of Statistical Software* 36, no. 3: 1–48. <https://doi.org/10.18637/jss.v036.i03>.
- Viechtbauer, W., and M. W.-L. Cheung. 2010. "Outlier and Influence Diagnostics for Meta-Analysis." *Research Synthesis Methods* 1, no. 2: 112–125. <https://doi.org/10.1002/jrsm.11>.
- Walkup, D. K., D. J. Leavitt, and L. A. Fitzgerald. 2017. "Effects of Habitat Fragmentation on Population Structure of Dune-Dwelling Lizards." *Ecosphere* 8, no. 3: 1729. <https://doi.org/10.1002/ecs2.1729>.
- Warton, D. I., R. A. Duursma, D. S. Falster, and S. Taskinen. 2012. "Smatr 3—An R Package for Estimation and Inference About Allometric Lines." *Methods in Ecology and Evolution* 3, no. 2: 257–259. <https://doi.org/10.1111/j.2041-210X.2011.00153.x>.
- Welsh, H. H., K. L. Pope, and C. A. Wheeler. 2008. "Using Multiple Metrics to Assess the Effects of Forest Succession on Population Status: A Comparative Study of Two Terrestrial Salamanders in the US Pacific Northwest." *Biological Conservation* 141: 14. <https://doi.org/10.1016/j.biocon.2008.02.014>.
- Williams, J. J., R. Freeman, F. Spooner, and T. Newbold. 2022. "Vertebrate Population Trends Are Influenced by Interactions Between Land Use, Climatic Position, Habitat Loss and Climate Change." *Global Change Biology* 28, no. 3: 797–815. <https://doi.org/10.1111/gcb.15978>.
- Data Sources**
- Abom, R., and L. Schwarzkopf. 2016. "Short-Term Responses of Reptile Assemblages to Fire in Native and Weedy Tropical Savannah." *Global Ecology and Conservation* 6: 58–66. <https://doi.org/10.1016/j.gecco.2016.02.002>.
- Craig, M. D., A. H. Grigg, M. J. Garkaklis, et al. 2009. "Does Habitat Structure Influence Capture Probabilities? A Study of Reptiles in a Eucalypt Forest." *Wildlife Research* 36, no. 6: 509. <https://doi.org/10.1071/WR09014>.
- Craig, M. D., A. H. Grigg, R. J. Hobbs, and G. E. S. J. Hardy. 2014. "Does Coarse Woody Debris Density and Volume Influence the Terrestrial Vertebrate Community in Restored Bauxite Mines?" *Forest Ecology and Management* 318: 142–150. <https://doi.org/10.1016/j.foreco.2014.01.011>.
- Craig, M. D., G. E. S. J. Hardy, J. B. Fontaine, et al. 2012. "Identifying Unidirectional and Dynamic Habitat Filters to Faunal Recolonisation in Restored Mine-Pits." *Journal of Applied Ecology* 49, no. 4: 919–928. <https://doi.org/10.1111/j.1365-2664.2012.02152.x>.
- Craig, M. D., R. J. Hobbs, A. H. Grigg, et al. 2010. "Do Thinning and Burning Sites Revegetated After Bauxite Mining Improve Habitat for Terrestrial Vertebrates?" *Restoration Ecology* 18, no. 3: 300–310. <https://doi.org/10.1111/j.1526-100X.2009.00526.x>.
- Craig, M. D., M. E. Smith, V. L. Stokes, G. E. S. J. Hardy, and R. J. Hobbs. 2018. "Temporal Longevity of Unidirectional and Dynamic Filters to Faunal Recolonization in Post-Mining Forest Restoration." *Austral Ecology* 43, no. 8: 973–988. <https://doi.org/10.1111/aec.12647>.
- Craig, M. D., V. L. Stokes, G. E. S. J. Hardy, and R. J. Hobbs. 2015. "Edge Effects Across Boundaries Between Natural and Restored Jarrah (*Eucalyptus marginata*) Forests in South-Western Australia." *Austral Ecology* 40, no. 2: 186–197. <https://doi.org/10.1111/aec.12193>.
- Davis, R. A., and M. D. Craig. 2024. "Long-Term Post-Fire Succession of Reptiles in an Urban Remnant in South-Western Australia." *International Journal of Wildland Fire* 33, no. 6: 24033. <https://doi.org/10.1071/WF24033>.
- Davis, R. A., and T. S. Doherty. 2015. "Rapid Recovery of an Urban Remnant Reptile Community Following Summer Wildfire." *PLoS One* 10, no. 5: e0127925. <https://doi.org/10.1371/journal.pone.0127925>.
- Doherty, T. S., D. F. Bohórquez Fandiño, D. J. Watchorn, S. M. Legge, and C. R. Dickman. 2024a. "Experimentally Testing Animal Responses to Prescribed Fire Size and Severity." *Conservation Biology* 38, no. 3: e14231. <https://doi.org/10.1111/cobi.14231>.
- Doherty, T. S., R. A. Davis, E. J. B. van Etten, N. Collier, and J. Krawiec. 2015. "Response of a Shrubland Mammal and Reptile Community to a History of Landscape-Scale Wildfire." *International Journal of Wildland Fire* 24, no. 4: 534. <https://doi.org/10.1071/WF14115>.
- Doherty, T. S., B. Johnson, G. R. Friend, and A. F. Wayne. 2024b. "Multi-Year Responses of Reptiles to Prescribed Burning in a Eucalypt Forest Ecosystem." *Austral Ecology* 49, no. 8: 13572. <https://doi.org/10.1111/aec.13572>.
- Dundas, S. J., K. X. Ruthrof, G. E. S. J. Hardy, and P. A. Fleming. 2021. "Some Like It Hot: Drought-Induced Forest Die-Off Influences Reptile Assemblages." *Acta Oecologica* 111: 103714. <https://doi.org/10.1016/j.actao.2021.103714>.
- Fitzgerald, L. A., W. A. Ryberg, D. K. Walkup, et al. 2022. "Ecology and Conservation of the Imperiled Dunes Sagebrush Lizard: An Ecological Specialist in a Threatened Ecosystem." In *Imperiled: The Encyclopedia of Conservation*, 494–510. Elsevier. <https://doi.org/10.1016/B978-0-12-821139-7.00214-2>.
- Hromada, S. J., C. A. F. Howey, M. B. Dickinson, R. W. Perry, W. M. Roosenburg, and C. M. Gienger. 2018. "Response of Reptile and Amphibian Communities to the Reintroduction of Fire in an Oak/Hickory Forest." *Forest Ecology and Management* 428: 1–13. <https://doi.org/10.1016/j.foreco.2018.06.018>.
- Irvine, C. 2024. *The Impacts of Urbanisation on Reptiles in a Young, Rapidly Urbanizing City*. University of Western Australia. <https://doi.org/10.26182/h4er-1v20>.
- Jessop, T. S., and G. R. Gillespie. 2022. "Monitoring Methods Influence Native Predator Detectability and Inferred Occupancy Responses to Introduced Carnivore Management." *Wildlife Research* 50, no. 1: 16–27. <https://doi.org/10.1071/WR22012>.
- Keehn, J. E., K. T. Shoemaker, and C. R. Feldman. 2019b. "Population-Level Effects of Wind Farms on a Desert Lizard." *Journal of Wildlife Management* 83, no. 1: 145–157. <https://doi.org/10.1002/jwmg.21565>.
- Lyndon-Gee, F., and T. S. Jessop. 2020. "Attenuated Phenotypic Responses of Lizard Morphology to Logging and Fire-Related Forest Disturbance." *Evolutionary Ecology* 34, no. 5: 841–853. <https://doi.org/10.1007/s10682-020-10067-9>.

- Macdonald, K. J., T. S. Doherty, B. A. Hradsky, and D. A. Driscoll. 2024b. "Disturbance and Ecosystem Management Interact to Shape Reptile Body Condition." *Animal Conservation* 28: 12983. <https://doi.org/10.1111/acv.12983>.
- Santos, X., J. Belliure, J. F. Gonçalves, and J. G. Pausas. 2022b. "Resilience of Reptiles to Megafires." *Ecological Applications* 32, no. 2: e2518. <https://doi.org/10.1002/eap.2518>.
- Smith, A. L. 2018b. "Successional Changes in Trophic Interactions Support a Mechanistic Model of Post-Fire Population Dynamics." *Oecologia* 186, no. 1: 129–139. <https://doi.org/10.1007/s00442-017-4016-z>.
- Smith, A. L., C. M. Bull, and D. A. Driscoll. 2012. "Post-Fire Succession Affects Abundance and Survival but Not Detectability in a Knob-Tailed Gecko." *Biological Conservation* 145, no. 1: 139–147. <https://doi.org/10.1016/j.biocon.2011.10.023>.
- Stokeld, D., A. Fisher, T. Gentles, et al. 2018. "Rapid Increase of Australian Tropical Savanna Reptile Abundance Following Exclusion of Feral Cats." *Biological Conservation* 225: 213–221. <https://doi.org/10.1016/j.biocon.2018.06.025>.
- Weiss, S. L., and R. M. Brower. 2021. "Wildfire as a Natural Stressor and Its Effect on Female Phenotype and Ornament Development." *Ecology and Evolution* 11, no. 11: 6223–6232. <https://doi.org/10.1002/ece3.7457>.
- Westaway, D. M., C. J. Jolly, D. R. Michael, and D. G. Nimmo. 2024. "Factors Affecting Microhabitat Use in Two Agamid Species From South-Eastern Australia." *Austral Ecology* 49: e13567. <https://doi.org/10.1111/aec.13567>.
- Woolley, C. K., S. Hartley, J. G. Innes, et al. 2023. "Conservation of Skinks in New Zealand Cities." *Urban Ecosystems* 26, no. 5: 1493–1508. <https://doi.org/10.1007/s11252-023-01398-w>.

Supporting Information

Additional supporting information can be found online in the Supporting Information section.