

THE DIRICHLET PROBLEM FOR THE 1-LAPLACIAN WITH A GENERAL SINGULAR TERM AND L^1 -DATA

MARTA LATORRE, FRANCESCANTONIO OLIVA, FRANCESCO PETITTA, AND SERGIO SEGURA DE LEÓN

ABSTRACT. We study the Dirichlet problem for an elliptic equation involving the 1-Laplace operator and a reaction term, namely:

$$\begin{cases} -\Delta_1 u = h(u)f(x) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where $\Omega \subset \mathbb{R}^N$ is an open bounded set having Lipschitz boundary, $f \in L^1(\Omega)$ is nonnegative, and h is a continuous real function that may possibly blow up at zero. We investigate optimal ranges for the data in order to obtain existence, nonexistence and (whenever expected) uniqueness of nonnegative solutions.

CONTENTS

1. Introduction	1
2. Preliminaries	3
2.1. Fixing some notation	3
2.2. Functions of bounded variation	4
2.3. L^∞ -divergence-measure vector fields	4
2.4. Construction of an auxiliary function	5
3. Assumptions and main results	6
4. Proof of the results of Section 3	7
4.1. Approximation scheme and a priori estimates	7
4.2. Estimates in BV and existence of an a.e. finite limit function u	9
4.3. Proof of the main existence result	10
4.4. Proof of the uniqueness result	15
5. The case of a nonnegative f	15
6. Remarks and examples	17
6.1. The case of a nonnegative h	17
6.2. The case $h(\infty) > 0$	17
6.3. Examples	18
Acknowledgement	20
References	20

1. INTRODUCTION

This paper is concerned with the Dirichlet problem for an equation involving the 1-Laplace operator and a lower order term in a bounded open set $\Omega \subset \mathbb{R}^N$ having Lipschitz boundary. More precisely, we deal with problem

$$\begin{cases} -\operatorname{div} \left(\frac{Du}{|Du|} \right) = h(u)f(x) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.1)$$

2010 *Mathematics Subject Classification.* 35J60, 35J75, 34B16, 35R99, 35A02.

Key words and phrases. 1-Laplacian, Nonlinear elliptic equations, Singular elliptic equations, L^1 data.

where $f \in L^1(\Omega)$ and h is a continuous real function; both are assumed to be nonnegative and we look for nonnegative solutions. The function $h(s)$ is not necessarily required to be bounded near $s = 0^+$.

Problems involving the 1-Laplace operator are known to be closely related to the mean curvature operator ([26]) and they enter in a variety of practical issues as (e.g. in the autonomous case $h \equiv 1$) in image restoration and in torsion problems ([16, 17, 27, 22, 5]). The nonautonomous/nonsingular case in relation with more theoretical subjects such as eigenvalues problems and critical Sobolev exponent has also been considered (see [14, 18, 8] and references therein). We refer the interested reader to the monograph [3] for a more complete review on applications in image processing.

For what concerns our study, in particular, it is worth recalling first the typical features of the related autonomous problem

$$-\operatorname{div} \left(\frac{Du}{|Du|} \right) = f(x) \quad \text{in } \Omega. \quad (1.2)$$

This problem has been usually addressed as the limit problem for p -laplacian type equations when p goes to 1^+ . In this setting it was considered by B. Kawohl in [16] for constant data and by M. Cicalese and C. Trombetti in [10] for data belonging to the Marcinkiewicz space $L^{N,\infty}(\Omega)$. General data belonging to $W^{-1,\infty}(\Omega)$, the dual space of $W_0^{1,1}(\Omega)$, are taken in [20]. Summarizing the results of these papers, it is proved that there exists a BV -function which is the limit of solutions to p -laplacian type equations only when the datum has small norm.

Existence of solution to (1.2) is also analyzed in [20] taken into account the notion of solution for equations involving the 1-Laplace operator introduced in [2, 14]. The L^1 -setting is studied in [21], where it is observed that if $f \in L^1(\Omega) \setminus W^{-1,\infty}(\Omega)$, no almost everywhere finite solution can be expected (see [21, Remarks 4.4 and 4.5]). More precisely, it is proved that to get an almost everywhere finite solution, the datum f must belong to $W^{-1,\infty}(\Omega)$ with $\|f\|_{W^{-1,\infty}(\Omega)} \leq 1$.

Concerning the presence of a non-constant term h in (1.1) one can refer to [18, 14, 8] for variational method in the study of eigenvalues type problems or critical Sobolev exponents. More recent papers analyzing the non autonomous case can be found in [15, 23]. The case of a, possibly singular, general nonlinearity h in (1.1) has been first addressed in [13] in the case $h(s) = s^{-\gamma}$ (with $0 < \gamma \leq 1$); here a BV -solution is found for every nonnegative $f \in L^N(\Omega)$. This existence result, without any smallness requirement, implies that the presence of $h(u)$ has a regularizing effect. Furthermore, in this paper it is shown the contrast between two cases of data: positive ($f > 0$) and nonnegative ($f \geq 0$). Roughly speaking, solutions of problem (1.1) with positive data have better properties and even a uniqueness result can be proved. In [12] (see also [24]) more general functions h and data f are analyzed and sharp existence and nonexistence results for (1.1) are given for nonnegative data in $f \in L^{N,\infty}(\Omega)$, completing the picture of [20] and, as we already mentioned, emphasizing the regularity effect given by the nonlinear term in order to get non-trivial solutions.

Here we want to go further focusing on a different type of regularizing effect given by the nonlinear term h if the data are assumed to be less regular. One of the purposes will in fact consist in discriminate whether a blow-up phenomenon as the one described in [21] does occur or not depending on the behavior of the nonlinearity at infinity. Our main aim then will be the identification of a general class of continuous functions h such that there exists a solution to (1.1) for every nonnegative $f \in L^1(\Omega)$. We prove that it is so when $h(s) \rightarrow 0$, as $s \rightarrow +\infty$. Moreover, if h attains the zero value, then we found a solution in $BV(\Omega) \cap L^\infty(\Omega)$, while if h is separated from zero (that is: $h(s) \geq m > 0$ for all $s > 0$), then existence of a BV -solution is only possible when $f \in W^{-1,\infty}(\Omega)$. Therefore, the closer to zero is h , the greater the regularizing effect. Also we address to the interplay between the behavior of h at infinity and the one near the origin, showing how this produces non-degenerate solutions no matter of the regularity or the smallness of the data.

The main difficulties in this type of analysis rely, on one hand, on well known issues related to the presence of the singular operator Δ_1 , as for instance the need to give sense to the singular quotient $|Du|^{-1}Du$, the invariance under monotone transformations that (e.g. in the autonomous case) leads to a structural non-uniqueness property. Moreover, one needs to work in the BV framework where weaker compactness results imply further complications as the ones related to the traces of the limit functions.

On the other hand, the presence of the general, and possibly singular, nonlinear term h makes the situation even worse as no monotonicity arguments apply and nearly all a priori estimates are only local. To get rid of these problems we first need to construct an auxiliary function that in some sense controls h and then to carefully estimate suitable truncations of the approximate solutions.

In order to pass to the limit then, one has to take care of the possibly singular behavior of the lower order term and this leads to develop an argument that allows to manage the set where the approximating solutions are small; this is done again by mean of suitably truncated auxiliary functions.

This paper is organized as follows. Section 2 is devoted to introduce some preliminary tools. In Section 3 we state our main assumptions, our notion of solution to problem (1.1) and our main results on existence and uniqueness in the case of positive data. These results are proved in Section 4. In Section 5, we analyze the case of nonnegative data. The last section is concerned with further remarks and examples also casting lights on the optimality of the presented results.

2. PRELIMINARIES

2.1. Fixing some notation. We denote by $\mathcal{H}^{N-1}(E)$ the $(N-1)$ -dimensional Hausdorff measure of a set E while $|E|$ stands for its N -dimensional Lebesgue measure. The characteristic function of a set E will be written as χ_E .

For the entire paper $\Omega \subset \mathbb{R}^N$ ($N \geq 1$) is open and bounded with Lipschitz boundary. Thus, an outward normal unit vector $\nu(x)$ is defined for \mathcal{H}^{N-1} -almost every $x \in \partial\Omega$. We will make use of the usual Lebesgue and Sobolev spaces, denoted by $L^q(\Omega)$ and $W_0^{1,p}(\Omega)$, respectively, and $C_c^1(\Omega)$ stands for the set of all functions with compact support which are continuously differentiable on Ω .

Positive functions, that is, those satisfying $f(x) > 0$ for almost all $x \in \Omega$, will be denoted by $f > 0$, while $f \geq 0$ stands for functions such that $f(x) \geq 0$ almost everywhere in Ω .

In what follows, $\mathcal{M}(\Omega)$ is the usual space of Radon measures with finite total variation over Ω . The space $\mathcal{M}_{\text{loc}}(\Omega)$ is the space of Radon measures which are locally finite in Ω , that is, measures with finite total variation on every $\omega \subset\subset \Omega$. Integrals with respect to Lebesgue measure will be denoted as $\int_{\Omega} f dx$, for a Lebesgue summable function f . For an integral with respect to a measure $\mu \in \mathcal{M}(\Omega)$ other than Lebesgue and Hausdorff ones we will use the notation $\int_{\Omega} f d\mu$, instead of $\int_{\Omega} f d\mu$.


For a fixed $k > 0$, the truncation function $T_k : \mathbb{R} \rightarrow \mathbb{R}$ is given by

$$T_k(s) := \max(-k, \min(s, k));$$

while, for each $\delta > 0$, we also use the following auxiliary function

$$V_{\delta}(s) := \begin{cases} 1 & s \leq \delta, \\ \frac{2\delta - s}{\delta} & \delta < s < 2\delta, \\ 0 & s \geq 2\delta. \end{cases} \quad (2.1)$$

We explicitly remark that, if no otherwise specified, we will denote by C several positive constants whose value may change from line to line and, sometimes, on the same line. These

FIGURE 1. Functions $T_k(s)$ and $V_\delta(s)$

values will only depend on the data but they will never depend on the indexes of the sequences we will introduce.

2.2. Functions of bounded variation. The space of functions of bounded variation is defined by

$$BV(\Omega) := \{u \in L^1(\Omega) : Du \in \mathcal{M}(\Omega)^N\}.$$

We also recall that every function $u \in BV(\Omega)$ has a trace defined on $\partial\Omega$ and $u|_{\partial\Omega} \in L^1(\partial\Omega)$. We underline that the $BV(\Omega)$ space, endowed with the norm

$$\|u\|_{BV(\Omega)} = \int_{\Omega} |u| dx + \int_{\Omega} |Du|,$$

or with

$$\|u\|_{BV(\Omega)} = \int_{\partial\Omega} |u| d\mathcal{H}^{N-1} + \int_{\Omega} |Du|,$$

is a Banach space and both norms are equivalent. We denote by $BV_{\text{loc}}(\Omega)$ the space of functions in $BV(\omega)$ for every $\omega \subset\subset \Omega$. By L_u we denote the set of approximate points of u , so that the approximate discontinuity set is $S_u = \Omega \setminus L_u$ and with J_u the set of the approximate jump points. It is well known that any function $u \in BV(\Omega)$ can be identified with its precise representative u^* , which is the Lebesgue representative \tilde{u} in L_u , while $u^* = \frac{u^+ + u^-}{2}$ in J_u where u^+, u^- are the one-sided limits of u . Moreover, it can be shown that $\mathcal{H}^{N-1}(S_u \setminus J_u) = 0$ and that u^* is \mathcal{H}^{N-1} -measurable. With an abuse of notation, in the sequel we will frequently write $g(u)^*$, where g is merely continuous and u belongs to the BV space, meaning that $g(u)^*$ is equal to $g(\tilde{u})$ in L_u while $g(u)^* = \frac{g(u^+) + g(u^-)}{2}$ in J_u .

When taken limits, we often apply the lower semicontinuity with respect to the L^1 -convergence of the functionals given by

$$\begin{aligned} u &\mapsto \int_{\Omega} \varphi |Du| && \text{if } \varphi \in C_0^1(\Omega) \text{ and } \varphi \geq 0, \\ u &\mapsto \int_{\Omega} |Du| + \int_{\partial\Omega} |u| \mathcal{H}^{N-1}. \end{aligned}$$

For a more complete review regarding BV spaces we refer to [1] from which we mainly derive our notation.

2.3. L^∞ -divergence-measure vector fields. We denote by

$$\mathcal{DM}^\infty(\Omega) := \{z \in L^\infty(\Omega)^N : \text{div } z \in \mathcal{M}(\Omega)\},$$

and by $\mathcal{DM}_{\text{loc}}^\infty(\Omega)$ its local version, namely: the space of bounded vector fields z such that $\text{div } z \in \mathcal{M}_{\text{loc}}(\Omega)$. For our purposes we recall the L^∞ -divergence-measure vector fields theory due to [4] and [9]. We first recall that if $z \in \mathcal{DM}^\infty(\Omega)$ then $\text{div } z$ is a measure absolutely continuous with respect to \mathcal{H}^{N-1} .

In [4] the following distribution $(z, Dv) : C_c^1(\Omega) \mapsto \mathbb{R}$ is considered:

$$\langle (z, Dv), \varphi \rangle := - \int_{\Omega} v^* \varphi \text{div } z - \int_{\Omega} v z \cdot \nabla \varphi dx. \quad (2.2)$$

In [21] and [7] the authors prove that (z, Dv) is well defined if $z \in \mathcal{DM}^\infty(\Omega)$ and $v \in BV(\Omega) \cap L^\infty(\Omega)$ since one can show that $v^* \in L^\infty(\Omega, \operatorname{div} z)$. Moreover, in [13] the authors show that (2.2) is well posed if $z \in \mathcal{DM}_{\operatorname{loc}}^\infty(\Omega)$ and $v \in BV_{\operatorname{loc}}(\Omega) \cap L^1_{\operatorname{loc}}(\Omega, \operatorname{div} z)$. In all cases, it holds that

$$|\langle (z, Dv), \varphi \rangle| \leq \|\varphi\|_{L^\infty(U)} \|z\|_{L^\infty(U)^N} \int_U |Dv|,$$

for all open set $U \subset\subset \Omega$ and for all $\varphi \in C_c^1(U)$. Moreover one has

$$\left| \int_B (z, Dv) \right| \leq \int_B |(z, Dv)| \leq \|z\|_{L^\infty(U)^N} \int_B |Dv|, \quad (2.3)$$

for all Borel sets B and for all open sets U such that $B \subset U \subset \Omega$.

We recall that in [4] it is proved that every $z \in \mathcal{DM}^\infty(\Omega)$ possesses a weak trace on $\partial\Omega$ of its normal component which is denoted by $[z, \nu]$ (recall that $\nu(x)$ is the outward normal unit vector). Moreover, it holds

$$\|[z, \nu]\|_{L^\infty(\partial\Omega)} \leq \|z\|_{L^\infty(\Omega)^N},$$

and it also satisfies that if $z \in \mathcal{DM}^\infty(\Omega)$ and $v \in BV(\Omega) \cap L^\infty(\Omega)$, then

$$v[z, \nu] = [vz, \nu], \quad (2.4)$$

(see [7]).

Finally, we will also use the Green formula due to [13]. If $z \in \mathcal{DM}_{\operatorname{loc}}^\infty(\Omega)$ and $v \in BV(\Omega) \cap L^\infty(\Omega)$ is such that $v^* \in L^1(\Omega, \operatorname{div} z)$, then $vz \in \mathcal{DM}^\infty(\Omega)$ and a weak trace can be defined as well as the following Green formula holds.

Lemma 2.1. *Let $z \in \mathcal{DM}_{\operatorname{loc}}^\infty(\Omega)$ and let $v \in BV(\Omega) \cap L^\infty(\Omega)$ be such that $v^* \in L^1(\Omega, \operatorname{div} z)$. Then $vz \in \mathcal{DM}^\infty(\Omega)$ and it holds*

$$\int_\Omega v^* \operatorname{div} z + \int_\Omega (z, Dv) = \int_{\partial\Omega} [vz, \nu] d\mathcal{H}^{N-1}. \quad (2.5)$$

2.4. Construction of an auxiliary function. Our method of finding a solution to problem (1.1) is based on the possibility of taking $1/h(u)$ as test function (at least when u is large enough). Obviously h is merely a continuous function and we are not allowed to use it. Hence, we need to have a suitable approximation \bar{h} of h which may be used. This subsection is devoted to define this approximation.

Let $h : (0, \infty) \rightarrow (0, \infty)$ be a continuous function such that there exists finite


$$\lim_{s \rightarrow \infty} h(s) := h(\infty).$$

Then, the aim is to construct a decreasing function $\bar{h} : (0, \infty) \rightarrow (0, \infty)$ satisfying

- a) $\bar{h}(s) \geq h(s)$ for all $s \geq 0$;
- b) \bar{h} is locally Lipschitz-continuous;
- c) $\lim_{s \rightarrow \infty} \bar{h}(s) := h(\infty)$.

We may define \bar{h} through the following stages.

- (i) In the first step, we get a function $h_1(s)$ which is a C^1 -approximation to $h(s) + e^{-s}$ lying between $h(s)$ and $h(s) + 2e^{-s}$. To this end, we proceed as in the proof of Meyers-Serrin's Theorem. Consider the intervals $(\frac{1}{n+2}, \frac{1}{n})$ and $(n, n+2)$, $n \in \mathbb{N}$, jointly with $(\frac{1}{2}, 2)$. They form an open cover of $(0, +\infty)$. Then a smooth partition of unity can be selected. Using mollifiers, choose a C^1 -approximation to $h(s) + e^{-s}$ between $h(s)$ and $h(s) + 2e^{-s}$ in each interval of the cover, and assemble them by means of the partition of unity to obtain a function h_1 . Thus, h_1 is a C^1 -function defined on $(0, +\infty)$ such that $h_1(s) \geq h(s)$ for all $s > 0$.

FIGURE 2. The construction of \bar{h}

- (ii) We denote by h_2 the rising sun function of h_1 (see, for instance, [28, Exercises 1G and 4F]). We recall that the set of all shadow points of h_1 is an open set and can be written as a disjoint union $\cup_{j \in \mathbb{N}} (a_j, b_j)$. The procedure to pass from h_1 to h_2 can easily be translated to their derivatives: It just consists in changing the value of h_1' on any (a_j, b_j) by defining $h_2'(s) = 0$. Observe that h_1' is a continuous function and so it is bounded from below on any interval $(0, n]$, with $n \in \mathbb{N}$. We infer that h_2' is non-positive and bounded from below on any interval $(0, n]$, with $n \in \mathbb{N}$, although it may be discontinuous at the points a_j . Hence, h_2 is a non-increasing locally Lipschitz-continuous function such that $h_2(s) \geq h(s)$ for all $s > 0$ and $\lim_{s \rightarrow \infty} h_2(s) := h(\infty)$.

- (iii) We define $\bar{h}(s) = h_2(s) + e^{-s}$ which is decreasing, and so a), b) and c) hold.

3. ASSUMPTIONS AND MAIN RESULTS

Consider the following Dirichlet problem

$$\begin{cases} -\Delta_1 u = h(u)f(x) & \text{in } \Omega, \\ u \geq 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (3.1)$$

where the function $h : [0, \infty) \mapsto (0, \infty]$ is continuous, finite outside the origin, and satisfying the following growth condition:

$$\exists c, s_1 > 0, \gamma \geq 0 \text{ such that } h(s) \leq \frac{c}{s^\gamma} \text{ if } s \leq s_1. \quad (\text{h1})$$

We also assume that $h(0) \neq 0$ and that there exists a finite limit at infinity of the function h , i.e.

$$\lim_{s \rightarrow \infty} h(s) := h(\infty). \quad (\text{h2})$$

Here we state the results relative to the case $h(\infty) = 0$, while the case $h(\infty) > 0$ will be discussed in Section 6.2 below.

The main feature of problem (3.1) is that the datum f is merely integrable. Here we state the result in case that f is positive, postponing the discussion of the nonnegative case to Section

5. Let us also underline that requiring that h is positive is not restrictive at all: we refer to Section 6.1 to handle the simpler case $h \geq 0$. Finally we point out that the case of a bounded h is also covered by the above assumptions.

It is useful to fix the following notation:

$$\sigma := \max(1, \gamma),$$

which will be used henceforth.

Definition 3.1. A nonnegative function u having $T_k(u) \in BV_{\text{loc}}(\Omega)$ for any $k > 0$ is a solution to problem (3.1) if there exists $z \in \mathcal{DM}^\infty(\Omega)$ with $\|z\|_{L^\infty(\Omega)^N} \leq 1$ such that $h(u)f \in L^1(\Omega)$ and

$$-\operatorname{div} z = h(u)f \quad \text{as measures in } \Omega, \quad (3.2)$$

$$(z, DT_k(u)) = |DT_k(u)| \quad \text{as measures in } \Omega \text{ for any } k > 0, \quad (3.3)$$

and one of the following conditions holds:

$$\lim_{\epsilon \rightarrow 0^+} \int_{\Omega \cap B(x, \epsilon)} T_k(u(y)) dy = 0 \quad \text{for any } k > 0 \quad \text{or} \quad [z, \nu](x) = -1 \quad \text{for } \mathcal{H}^{N-1}\text{-a.e. } x \in \partial\Omega. \quad (3.4)$$

Remark 3.2. Let us observe that Definition 3.1 extends the ones in [13, 12] the main difference being that, as no BV solutions (nor bounded one) are expected in general (see Example 1 in Section 6.3), then conditions (3.3) and (3.4) have to be satisfied by truncations of the solution. Moreover it is also directly related to the definition given in [21] in the *infinite energy* regime of the autonomous case. A final comment on the weak way the boundary datum is assumed; if $T_k(u)$ does possesses a trace in the usual BV sense (e.g. if $\gamma \leq 1$, as we shall see) then the Dirichlet boundary condition (3.4) reduces to the more easily readable one

$$T_k(u)(1 + [z, \nu](x)) = 0 \quad \text{for } \mathcal{H}^{N-1}\text{-a.e. } x \in \partial\Omega.$$

We now state existence and uniqueness theorems in the case $f > 0$:

Theorem 3.3. Let $0 < f \in L^1(\Omega)$ and let h satisfy (h1) and (h2) with $h(\infty) = 0$. Then there exists a solution u to problem (3.1) in the sense of Definition 3.1 which is almost everywhere finite.

Theorem 3.4. Let $0 < f \in L^1(\Omega)$ and let h be decreasing. Then there exists at most one solution u to (3.1) in the sense of Definition 3.1 in the class $T_k^\sigma(u) \in BV(\Omega)$ for any $k > 0$.

We also have the following results that fit with the known nonexistence ones and that highlight the regularization effect produced by the nonlinear term:

Proposition 3.5. If u is a solution to (3.1), then $h(u)f \in W^{-1, \infty}(\Omega)$ and $\|h(u)f\|_{W^{-1, \infty}(\Omega)} \leq 1$.

Proposition 3.6. Let u be the solution to (3.1) found in Theorem 3.3. Then it holds that $T_k^\sigma(u) \in BV(\Omega)$ for any $k > 0$. Moreover if $h(0) = \infty$ then $u > 0$ almost everywhere in Ω .

4. PROOF OF THE RESULTS OF SECTION 3

4.1. Approximation scheme and a priori estimates. For $p > 1$, we consider the following scheme of approximation

$$\begin{cases} -\Delta_p u_p = h_p(u_p) f_p(x) & \text{in } \Omega, \\ u_p = 0 & \text{on } \partial\Omega, \end{cases} \quad (4.1)$$

where $h_p(s) := T_{\frac{1}{p-1}}(h(s))$ and $f_p = T_{\frac{1}{p-1}}(f)$. The existence of a nonnegative weak solution $u_p \in W_0^{1, p}(\Omega) \cap L^\infty(\Omega)$ is guaranteed by [19]. We highlight that, since we are taking $p \rightarrow 1^+$, without loss of generality we can assume that $1 < p < 2$. The a priori estimates on the approximate solutions do not depend of the sign of the datum f so that they shall be established for general nonnegative f 's and they'll also be used in Section 5.

For a reason that will be clear in a moment, we consider the function \bar{h} constructed in Section 2.4 relative to $g(s) = T_l(h(s))$ and $l = \sup_{s \in [s_1, \infty)} h(s)$ (i.e. we cut h only near zero) and we set $s_2 >$

$\max\{1, s_1\}$ satisfying $\bar{h}(s_2) < s_1^{-\sigma}$; the function \bar{h} is locally Lipschitz-continuous and satisfies $\bar{h}(s) \geq h(s)$ for any $s \geq s_2$ and $\lim_{s \rightarrow \infty} \bar{h}(s) = \lim_{s \rightarrow \infty} h(s)$. In order to get a first basic estimate we need a test function that mimics $\frac{1}{h(u)}$, to this end we define

$$\Phi(s) := \begin{cases} s^\sigma, & s < s_1, \\ \Psi(s), & s_1 \leq s \leq s_2, \\ 1/\bar{h}(s), & s > s_2, \end{cases}$$

where Ψ is a Lipschitz increasing function which makes Φ locally Lipschitz as well. Moreover we denote by

$$\Gamma_p(s) := \int_0^s \Phi'(t)^{\frac{1}{p}} dt.$$

We have the following easy but fundamental a priori estimate:

Lemma 4.1. *Let h satisfy (h1) and (h2) and let $f \in L^1(\Omega)$ be nonnegative. Let u_p be a solution of (4.1) then*

$$\|\Gamma_p(u_p)\|_{W_0^{1,p}(\Omega)} \leq C, \quad (4.2)$$

for some constant C which does not depend on p .

Proof. Recalling that $\sigma \geq 1$, one observes that $\Gamma_p(0) = 0$ and that u_p has zero trace, so that $\Gamma_p(u_p) \in W_0^{1,p}(\Omega)$.

Let us take $\Phi(u_p)$ as a test function in (4.1) and use the properties of \bar{h} , obtaining

$$\begin{aligned} \int_{\Omega} |\nabla \Gamma_p(u_p)|^p dx &= \int_{\Omega} |\nabla u_p|^p \Phi'(u_p) dx = \int_{\Omega} h_p(u_p) f_p \Phi(u_p) dx \\ &\leq c \int_{\{u_p \leq s_1\}} f dx + \left[\max_{s \in (s_1, s_2)} h(s) \right] \Phi(s_2) \int_{\{s_1 < u_p < s_2\}} f dx + \int_{\{u_p \geq s_2\}} f dx \leq C, \end{aligned}$$

for some positive constant C which does not depend on p . \square

Here is the second *global* a priori estimate that involves the truncations of u_p :

Lemma 4.2. *Let h satisfy (h1) and (h2), $f \in L^1(\Omega)$ be nonnegative, and let us fix $k > 0$. Then u_p satisfies*

$$\|T_k^{\frac{\sigma-1+p}{p}}(u_p)\|_{W_0^{1,p}(\Omega)} \leq C_k, \quad (4.3)$$

for some constant C_k which does not depend on p .

Proof. Let us take $T_k^\sigma(u_p)$ as a test function in (4.1) yielding to

$$\begin{aligned} \sigma \left(\frac{p}{\sigma-1+p} \right)^p \int_{\Omega} |\nabla T_k^{\frac{\sigma-1+p}{p}}(u_p)|^p dx &= \int_{\Omega} h_p(u_p) f_p T_k^\sigma(u_p) dx \\ &\leq c \int_{\{u_p \leq s_1\}} f dx + \left[\sup_{s \in [s_1, \infty)} h(s) \right] k^\sigma \int_{\{u_p > s_1\}} f dx \leq C_k, \end{aligned}$$

for some positive constant C_k which does not depend on p . \square

As we saw if $\gamma > 1$ then only a power of the truncations of u_p is bounded globally; this is a well known phenomenon in singular type problems (see e.g. [6, 25]). More in general we have the following local estimates

Lemma 4.3. *Let h satisfy (h1) and (h2) and let $f \in L^1(\Omega)$ be nonnegative. Then for any $\omega \subset\subset \Omega$*

$$\|T_k(u_p)\|_{W^{1,p}(\omega)} \leq C_\omega, \quad (4.4)$$

for some constant C_ω which does not depend on p .

Proof. Let $\varphi \in C_c^1(\Omega)$ be a nonnegative function with compact support in Ω such that $\varphi = 1$ in $\omega \subset\subset \Omega$; we take $(T_k(u_p) - k)\varphi^p$ as a test function in (4.1), in order to get

$$\int_{\Omega} |\nabla T_k(u_p)|^p \varphi^p dx + p \int_{\Omega} |\nabla T_k(u_p)|^{p-2} \nabla T_k(u_p) \cdot \nabla \varphi \varphi^{p-1} (T_k(u_p) - k) dx \leq 0.$$

After applying twice the Young inequality, we have

$$\begin{aligned} \int_{\Omega} |\nabla T_k(u_p)|^p \varphi^p dx &\leq (p-1)k\varepsilon \int_{\Omega} |\nabla T_k(u_p)|^p \varphi^p dx + k\varepsilon^{1-p} \int_{\Omega} |\nabla \varphi|^p dx \\ &\leq k\varepsilon \int_{\Omega} |\nabla T_k(u_p)|^p \varphi^p dx + k\varepsilon^{1-p} \int_{\Omega} |\nabla \varphi|^2 dx + k\varepsilon^{1-p} |\Omega|, \end{aligned}$$

and taking ε sufficiently small one easily concludes. \square

4.2. Estimates in BV and existence of an a.e. finite limit function u . The estimates found in Lemmas 4.1, 4.2 and 4.3 translate into uniform estimates in BV with respect to p that we collect in the following:

Lemma 4.4. *Let h satisfy (h1) and (h2) and let $f \in L^1(\Omega)$ nonnegative. Let u_p be a solution of (4.1) then it holds that*

$$\|\Gamma_p(u_p)\|_{BV(\Omega)} \leq C, \quad (4.5)$$

$$\|T_k^{\frac{\sigma-1+p}{p}}(u_p)\|_{BV(\Omega)} \leq C_k, \quad (4.6)$$

and for any $\omega \subset\subset \Omega$

$$\|T_k(u_p)\|_{BV(\omega)} \leq C_{\omega}, \quad (4.7)$$

for some constants C, C_k and C_{ω} which do not depend on p .

Proof. The proofs of (4.5), (4.6) and (4.7) are identical and rely on an application of the Young inequality. In fact, one has,

$$\int_{\Omega} |\nabla \Gamma_p(u_p)| dx \leq \int_{\Omega} |\nabla \Gamma_p(u_p)|^p dx + |\Omega| \stackrel{(4.2)}{\leq} C + |\Omega|,$$

which gives estimates (4.5) since the trace of u_p is zero and $\Gamma(0) = 0$. Similarly,

$$\int_{\Omega} |\nabla T_k^{\frac{\sigma-1+p}{p}}(u_p)| dx \leq \int_{\Omega} |\nabla T_k^{\frac{\sigma-1+p}{p}}(u_p)|^p dx + |\Omega| \stackrel{(4.3)}{\leq} C_k + |\Omega|,$$

and, if $\omega \subset\subset \Omega$, one has

$$\int_{\omega} |\nabla T_k(u_p)| dx \leq \int_{\omega} |\nabla T_k(u_p)|^p dx + |\omega| \stackrel{(4.4)}{\leq} C_{\omega} + |\omega|.$$

\square

We then deduce the following compactness results in which the most remarkable feature is given by the fact that the limit u is finite a.e. on Ω . As mentioned, this is due to the fact that h vanishes at infinity and it is one of the striking differences with respect to the autonomous case considered in [21].

Corollary 4.5. *Let h satisfy (h1) and (h2) with $h(\infty) = 0$ and let $f \in L^1(\Omega)$ be a nonnegative function. Let u_p be a solution of (4.1) then there exists a function u almost everywhere finite such that, up to subsequences, u_p converges to u almost everywhere in Ω as $p \rightarrow 1^+$. Moreover, one has that*

- i) $\Gamma_p(u_p)$ converges to $\Phi(u)$ in $L^q(\Omega)$ with $1 \leq q < \frac{N}{N-1}$ and $\nabla \Gamma_p(u_p)$ converges *-weakly as measures to $D\Phi(u)$;
- ii) $T_k^{\frac{\sigma-1+p}{p}}(u_p)$ converges to $T_k^{\sigma}(u)$ in $L^q(\Omega)$ with $1 \leq q < \frac{N}{N-1}$ and $\nabla T_k^{\frac{\sigma-1+p}{p}}(u_p)$ converges *-weakly as measures to $DT_k^{\sigma}(u)$ for any $k > 0$;
- iii) $T_k(u_p)$ locally converges to $T_k(u)$ in $L^q(\Omega)$ with $1 \leq q < \frac{N}{N-1}$ and $\nabla T_k(u_p)$ converges locally *-weakly as measures to $DT_k(u)$ for any $k > 0$.

Proof. By compactness in BV it follows from Lemma 4.4 that, up to subsequences, $v_p := \Gamma_p(u_p)$ converges to a function $v \in BV(\Omega)$ a.e. and in $L^q(\Omega)$ for every $1 \leq q < \frac{N}{N-1}$ and ∇v_p converges *-weakly as measures to Dv . Observe that if we define $u := \Phi^{-1}(v)$ then $u_p = \Gamma_p^{-1}(v_p)$ converges almost everywhere as $p \rightarrow 1^+$ to u . Finally, as $v = \Phi(u) \in L^q(\Omega)$ for all $1 \leq q < \frac{N}{N-1}$, one has that u is almost everywhere finite since v is almost everywhere finite and $\Phi(s)$ blows up only as $s \rightarrow \infty$. This shows that u_p converges almost everywhere to u as $p \rightarrow 1^+$ and it also proves i). The proofs of ii) and iii) follow in a similar way. \square

4.3. Proof of the main existence result. We will divide the proof in different lemmas, and then the proof of the existence will follow as the collection of these lemmas. First of all, let us underline that, from here on (even when not specified), u_p will be a solution to (4.1) and u is the almost everywhere limit as $p \rightarrow 1^+$ given by Corollary 4.5. The first result concerns the existence of the vector field z playing the role of the singular quotient $|Du|^{-1}Du$. In particular, the following lemma is the only place where the positivity of f in Ω will be employed.

Lemma 4.6. *Let h satisfy (h1) and (h2) provided $h(\infty) = 0$, and let $0 < f \in L^1(\Omega)$. Let u_p be a solution of (4.1) then there exists $z \in \mathcal{DM}^\infty(\Omega)$ with $\|z\|_{L^\infty(\Omega)^N} \leq 1$ such that $h(u)f \in L^1(\Omega)$ and such that*

$$-\operatorname{div} z = h(u)f \text{ as measures in } \Omega. \quad (4.8)$$

Moreover,

$$(z, DT_k(u)) = |DT_k(u)| \quad \text{as measures in } \Omega \text{ for any } k > 0. \quad (4.9)$$

Proof. We begin by considering the set \mathcal{N} of all levels $k > 0$ such that $|\{u = k\}| > 0$, which is a countable set. We point out that if $k \notin \mathcal{N}$, then

$$\chi_{\{u_p < k\}} \rightarrow \chi_{\{u < k\}} \quad \text{strongly in } L^r(\Omega) \text{ for all } 1 \leq r < \infty,$$

owing to the pointwise convergence $u_p \rightarrow u$ a.e. in Ω .

For a fixed $\omega \subset\subset \Omega$ one obviously has

$$\chi_{\omega \cap \{u_p < k\}} \rightarrow \chi_{\omega \cap \{u < k\}} \quad \text{strongly in } L^r(\omega) \text{ for all } 1 \leq r < \infty. \quad (4.10)$$

On the other hand, applying (4.4) and the Hölder inequality we deduce that for $1 \leq q < \frac{p}{p-1}$ and for any $k > 0$,

$$\| |\nabla T_k(u_p)|^{p-2} \nabla T_k(u_p) \|_{L^q(\omega)^N} \leq \left(\int_\omega |\nabla T_k(u_p)|^p dx \right)^{\frac{p-1}{p}} |\omega|^{\frac{1}{q} - \frac{p-1}{p}} \leq C_\omega^{\frac{p-1}{p}} |\omega|^{\frac{1}{q} - \frac{p-1}{p}}, \quad (4.11)$$

that is, the family $((|\nabla T_k(u_p)|^{p-2} \nabla T_k(u_p)) \chi_\omega)_p$ is bounded in $L^q(\omega)^N$ for every $1 \leq q < \infty$. This implies the existence of a vector field $z_{q,k} \in L^q(\omega)^N$ such that, up to subsequences,

$$(|\nabla T_k(u_p)|^{p-2} \nabla T_k(u_p)) \chi_\omega \rightharpoonup z_{q,k} \quad \text{weakly in } L^q(\omega)^N.$$

Now a standard diagonal argument takes to the existence of a unique vector field z_k , defined independently of q , such that

$$(|\nabla T_k(u_p)|^{p-2} \nabla T_k(u_p)) \chi_\omega \rightharpoonup z_k \quad \text{weakly in } L^q(\omega)^N \text{ for any } 1 \leq q < \infty. \quad (4.12)$$

Moreover, letting $p \rightarrow 1^+$, by weakly lower semicontinuity applied to (4.11), one yields to $\|z_k\|_{L^q(\omega)^N} \leq |\omega|^{\frac{1}{q}}$ for any $q < \infty$ and letting $q \rightarrow \infty$ we also have $\|z_k\|_{L^\infty(\omega)^N} \leq 1$.

The vector fields we have considered are defined in ω : in order to explicit this dependence, we will write z_k^ω for $k \notin \mathcal{N}$. It remains to extend them to the whole Ω . To this end, just take into account (4.12) to see that $\omega_1 \subset \omega_2 \subset\subset \Omega$ implies $z_k^{\omega_2}|_{\omega_1} = z_k^{\omega_1}$. Thus, it is enough to take an increasing sequence $\omega_n \subset\subset \Omega$ satisfying $\cup_{n=1}^\infty \omega_n = \Omega$ and so we obtain vector fields z_k ($k \notin \mathcal{N}$) such that $z_k|_\omega = z_k^\omega$ and (4.12) holds for each $\omega \subset\subset \Omega$. It also follows from $\|z_k^{\omega_n}\|_{L^\infty(\omega_n)^N} \leq 1$ for all $n \in \mathbb{N}$ that $\|z_k\|_{L^\infty(\Omega)^N} \leq 1$.

Now, choose $h, k \notin \mathcal{N}$ satisfying $0 < k < h$. Having in mind (4.12) and (4.10), and letting p go to 1 in the identity

$$(|\nabla T_h(u_p)|^{p-2} \nabla T_h(u_p)) \chi_{\omega \cap \{u_p < k\}} = (|\nabla T_k(u_p)|^{p-2} \nabla T_k(u_p)) \chi_{\omega},$$

we obtain

$$z_h \chi_{\{u < k\}} = z_k. \quad (4.13)$$

As (4.13) is in force, one can define the vector field z in Ω as

$$z(x) = z_k(x) \quad \text{when } u(x) < k \quad (k \notin \mathcal{N}).$$

It is straightforward that this vector field z is well-defined in Ω (since u is almost everywhere finite), $z \in L^\infty(\Omega)^N$ and $\|z\|_{L^\infty(\Omega)^N} \leq 1$. Furthermore,

$$z \chi_{\{u < k\}} = z_k \quad \text{whatever } k \notin \mathcal{N}.$$

Now we look at (4.8); let us denote

$$S_n(s) = \begin{cases} 1 & \text{if } s \leq n, \\ -s + n + 1 & \text{if } n < s < n + 1, \\ 0 & \text{if } s \geq n + 1, \end{cases}$$

and let $0 \leq \varphi \in C_c^1(\Omega)$; we take $S_n(u_p)\varphi$ as a test function in (4.1), yielding to

$$\int_{\Omega} |\nabla u_p|^{p-2} \nabla u_p \cdot \nabla \varphi S_n(u_p) dx - \int_{\{n < u_p < n+1\}} |\nabla u_p|^p \varphi dx = \int_{\Omega} h_p(u_p) f_p S_n(u_p) \varphi dx, \quad (4.14)$$

whence getting rid of the second term on the left hand side of (4.14) and taking $p \rightarrow 1^+$ one has

$$\int_{\Omega} z \cdot \nabla \varphi S_n(u) dx \geq \int_{\Omega} h(u) f S_n(u) \varphi dx,$$

where, on the right hand side, we have applied the Fatou Lemma. Again thanks to Fatou's Lemma we take $n \rightarrow \infty$, getting

$$\int_{\Omega} z \cdot \nabla \varphi dx \geq \int_{\Omega} h(u) f \varphi dx,$$

from which one deduces that $z \in \mathcal{DM}_{\text{loc}}^\infty(\Omega)$ and $h(u)f \in L_{\text{loc}}^1(\Omega)$. Let also observe that, in case $h(0) = \infty$, having $h(u)f \in L_{\text{loc}}^1(\Omega)$ implies that

$$\{u = 0\} \subset \{f = 0\}, \quad (4.15)$$

up to a set of zero Lebesgue measure; in particular, as $f > 0$ one has that $u > 0$. In order to prove that (4.8) holds we now consider $1 - S_n(u_p)$ as a test function in (4.1) that yields

$$\int_{\{n < u_p < n+1\}} |\nabla u_p|^p dx = \int_{\Omega} h_p(u_p) f_p (1 - S_n(u_p)) dx,$$

and taking the limit as $p \rightarrow 1^+$ by means of the dominated convergence Theorem (observe that $u_p \geq n$), one gets

$$\lim_{p \rightarrow 1^+} \int_{\{n < u_p < n+1\}} |\nabla u_p|^p dx = \int_{\Omega} h(u) f (1 - S_n(u)) dx,$$

and as $n \rightarrow \infty$, we have

$$\lim_{n \rightarrow \infty} \lim_{p \rightarrow 1^+} \int_{\{n < u_p < n+1\}} |\nabla u_p|^p dx = 0, \quad (4.16)$$

where in the last step we used again dominated convergence since

$$h(u) f \chi_{\{n < u < n+1\}} \leq \sup_{s \geq 1} h(s) f \in L^1(\Omega).$$

Now we want to pass to the limit every terms of (4.14) first in p and then in n . We easily pass to the limit the first term as well as the second term which goes to zero thanks to (4.16).

Concerning the term on the right hand side the limit is easy if $h(0) < \infty$. Hence assume $h(0) = \infty$. Let $\delta > 0$ such that $\delta \notin \mathcal{N}$, and write

$$\int_{\Omega} h_p(u_p) f_p S_n(u_p) \varphi dx = \int_{\{u_p < \delta\}} h_p(u_p) f_p S_n(u_p) \varphi dx + \int_{\{u_p \geq \delta\}} h_p(u_p) f_p S_n(u_p) \varphi dx; \quad (4.17)$$

we want to pass to the limit in the following order: p , then n , and finally $\delta \rightarrow 0^+$. We first pass to the limit the second term on the right hand side of (4.17); we observe that

$$h_p(u_p) f_p S_n(u_p) \varphi \chi_{\{u_p > \delta\}} \leq \sup_{s \in [\delta, \infty)} h(s) f \varphi \in L^1(\Omega),$$

which allows us to apply Lebesgue's Theorem as $p \rightarrow 1^+$. One gains that

$$\lim_{p \rightarrow 1^+} \int_{\{u_p \geq \delta\}} h_p(u_p) f_p S_n(u_p) \varphi dx = \int_{\{u \geq \delta\}} h(u) f S_n(u) \varphi dx,$$

and observing that $h(u) f \in L^1_{\text{loc}}(\Omega)$, then one can apply once again the Lebesgue Theorem first in n and then in δ . Hence, one has

$$\lim_{\delta \rightarrow 0^+} \lim_{n \rightarrow \infty} \lim_{p \rightarrow 1^+} \int_{\{u_p \geq \delta\}} h_p(u_p) f_p S_n(u_p) \varphi dx = \int_{\{u > 0\}} h(u) f \varphi dx. \quad (4.18)$$

Therefore, we are just left to show that the first term on the right hand side of (4.17) vanishes. To this aim we take $V_\delta(u_p) \varphi$ ($0 \leq \varphi \in C_c^1(\Omega)$ and $V_\delta(s)$ is defined in (2.1)) as a test function in (4.1) yielding to

$$\begin{aligned} \int_{\{u_p < \delta\}} h_p(u_p) f_p \varphi dx &\leq \int_{\Omega} h_p(u_p) f_p V_\delta(u_p) \varphi dx \\ &= \int_{\Omega} |\nabla u_p|^{p-2} \nabla u_p \cdot \nabla \varphi V_\delta(u_p) dx - \frac{1}{\delta} \int_{\{\delta < u_p < 2\delta\}} |\nabla u_p|^p \varphi dx, \end{aligned}$$

which gives

$$\int_{\{u_p < \delta\}} h_p(u_p) f_p \varphi dx \leq \int_{\Omega} |\nabla u_p|^{p-2} \nabla u_p \cdot \nabla \varphi V_\delta(u_p) dx. \quad (4.19)$$

Hence we can take $p \rightarrow 1^+$ in (4.19) obtaining

$$\limsup_{p \rightarrow 1^+} \int_{\{u_p < \delta\}} h_p(u_p) f_p \varphi dx \leq \int_{\Omega} z \cdot \nabla \varphi V_\delta(u) dx;$$

as there is no dependence on n , we let $\delta \rightarrow 0^+$ obtaining

$$\lim_{\delta \rightarrow 0^+} \limsup_{p \rightarrow 1^+} \int_{\{u_p < \delta\}} h_p(u_p) f_p \varphi dx \leq \int_{\{u=0\}} z \cdot \nabla \varphi dx \stackrel{(4.15)}{=} 0. \quad (4.20)$$

Summing up (recall that $S_n(s) \leq 1$) (4.20) implies that

$$\lim_{\delta \rightarrow 0^+} \lim_{n \rightarrow \infty} \lim_{p \rightarrow 1^+} \int_{\{u_p < \delta\}} h_p(u_p) f_p S_n(u_p) \varphi dx = 0,$$

which jointly with (4.18) gives

$$\lim_{n \rightarrow \infty} \lim_{p \rightarrow 1^+} \int_{\Omega} h_p(u_p) f_p S_n(u_p) \varphi dx = \int_{\{u > 0\}} h(u) f \varphi dx \stackrel{(4.15)}{=} \int_{\Omega} h(u) f \varphi dx.$$

This proves that (4.8) holds. Let us also underline that the validity of (4.8) allows us to apply Lemma 5.3 of [12] in order to deduce that $\text{div } z \in L^1(\Omega)$, i.e. both $z \in \mathcal{DM}^\infty(\Omega)$ and $h(u) f \in L^1(\Omega)$. In order to prove (4.9) we first need to show that a slightly modified version of (4.8) holds. For $k > 0$ we consider $(\rho_\epsilon * T_k^\sigma(u)) \varphi$ as a test function in (4.8) where ρ_ϵ is a sequence of standard mollifiers and $\varphi \in C_c^1(\Omega)$ is nonnegative. Hence

$$- \int_{\Omega} (\rho_\epsilon * T_k^\sigma(u)) \varphi \text{div } z = \int_{\Omega} h(u) f (\rho_\epsilon * T_k^\sigma(u)) \varphi dx.$$

Observe that $T_k^\sigma(u) \in BV(\Omega)$, that $(\rho_\epsilon * T_k^\sigma(u)) \rightarrow T_k^\sigma(u)^* \mathcal{H}^{N-1}$ -almost everywhere as $\epsilon \rightarrow 0^+$, and that $\rho_\epsilon * T_k^\sigma(u) \leq k^\sigma$. Then letting $\epsilon \rightarrow 0^+$ one has

$$-T_k^\sigma(u)^* \operatorname{div} z = h(u) f T_k^\sigma(u) \text{ as measures in } \Omega. \quad (4.21)$$

Now we take $T_k^\sigma(u_p) S_n(u_p) \varphi$ ($0 \leq \varphi \in C_c^1(\Omega)$ and $n > k$) as a test function in (4.1), deducing, after an application of the Young inequality, that

$$\begin{aligned} & \int_{\Omega} |\nabla T_k^{\frac{\sigma-1+p}{p}}(u_p)| \varphi dx + \int_{\Omega} T_k^\sigma(u_p) S_n(u_p) |\nabla u_p|^{p-2} \nabla u_p \cdot \nabla \varphi dx - k^\sigma \int_{\{n < u_p < n+1\}} |\nabla u_p|^p \varphi dx \\ & \leq \int_{\Omega} h_p(u_p) f_p T_k^\sigma(u_p) S_n(u_p) \varphi dx + \frac{p-1}{p} \left(\frac{p}{\sigma-1+p} \right)^{\frac{p^2}{p-1}} \int_{\Omega} \varphi dx, \end{aligned} \quad (4.22)$$

and we want to let $p \rightarrow 1^+$ first, and then $n \rightarrow \infty$ in the (4.22). We observe that by (4.16) the third term on the left hand side of (4.22) goes to zero as $p \rightarrow 1^+$ and $n \rightarrow \infty$. Then by lower semicontinuity and weak convergence at the left hand side of (4.22), yielding to

$$\begin{aligned} & \int_{\Omega} \varphi |DT_k^\sigma(u)| + \int_{\Omega} T_k^\sigma(u) S_n(u) z \cdot \nabla \varphi dx - \lim_{p \rightarrow 1^+} k^\sigma \int_{\{n < u_p < n+1\}} |\nabla u_p|^p \varphi dx \\ & \leq \int_{\Omega} h(u) f T_k^\sigma(u) S_n(u) \varphi dx, \end{aligned}$$

where in order to pass to the limit on the first term of right hand side we used dominated convergence Theorem as

$$h_p(u_p) f_p T_k^\sigma(u_p) S_n(u_p) \varphi \leq c f \varphi \chi_{\{u_p \leq s_1\}} + \max_{s \in [s_1, \infty)} h(s) k^\sigma f \varphi \chi_{\{u_p > s_1\}} \leq (c + \max_{s \in [s_1, \infty)} h(s) k^\sigma) f \varphi.$$

Finally one can take $n \rightarrow \infty$, obtaining

$$\int_{\Omega} \varphi |DT_k^\sigma(u)| + \int_{\Omega} T_k^\sigma(u) z \cdot \nabla \varphi dx \leq \int_{\Omega} h(u) f T_k^\sigma(u) \varphi dx \stackrel{(4.21)}{=} - \int_{\Omega} T_k^\sigma(u)^* \varphi \operatorname{div} z.$$

Recalling (2.2) the previous implies that

$$\int_{\Omega} \varphi |DT_k^\sigma(u)| \leq \int_{\Omega} \varphi (z, DT_k^\sigma(u)), \quad \forall \varphi \in C_c^1(\Omega), \quad \varphi \geq 0,$$

and since the reverse inequality is trivial, one has that

$$(z, DT_k^\sigma(u)) = |DT_k^\sigma(u)| \text{ as measures in } \Omega, \quad (4.23)$$

for any $k > 0$. Finally one can apply Proposition 4.5 of [11] which implies

$$\theta(z, DT_k(u), x) = \theta(z, DT_k^\sigma(u), x) \quad \text{for } |DT_k(u)|\text{-a.e. } x \in \Omega, \quad (4.24)$$

where $\theta(z, DT_k(u), \cdot)$ is the Radon-Nikodým derivative of the measure $(z, DT_k(u))$ with respect to $|DT_k(u)|$, while $\theta(z, DT_k^\sigma(u), \cdot)$ denotes the Radon-Nikodým derivative of $(z, DT_k^\sigma(u))$ with respect to $|DT_k^\sigma(u)|$. This, jointly with (4.23), implies (4.9). This concludes the proof of the lemma. \square

Now we show that the Dirichlet boundary condition holds.

Lemma 4.7. *Let h satisfy (h1) and (h2) with $h(\infty) = 0$ and let $0 < f \in L^1(\Omega)$. Let u be the function found in Lemma 4.5 and let z be the vector field of Lemma 4.6, then it holds*

$$\lim_{\epsilon \rightarrow 0^+} \int_{\Omega \cap B(x, \epsilon)} T_k(u(y)) dy = 0 \quad \text{or} \quad [z, \nu](x) = -1 \quad \text{for } \mathcal{H}^{N-1}\text{-a.e. } x \in \partial\Omega, \quad (4.25)$$

for any $k > 0$.

Proof. First of all we take $T_k^\sigma(u_p)$ as a test function in (4.1) and we apply the Young inequality in order to obtain (recalling that the trace of u_p is zero)

$$\int_{\Omega} |\nabla T_k^{\frac{\sigma-1+p}{p}}(u_p)| dx + \int_{\partial\Omega} T_k^{\frac{\sigma-1+p}{p}}(u_p) d\mathcal{H}^{N-1} \leq \int_{\Omega} h_p(u_p) f_p T_k^\sigma(u_p) dx + \frac{p-1}{p} \left(\frac{p}{\sigma-1+p} \right)^{\frac{p^2}{p-1}} |\Omega|.$$

By lower semicontinuity on the left hand side and by Lebesgue's dominated convergence Theorem on the right hand side, letting $p \rightarrow 1^+$, one has

$$\int_{\Omega} |DT_k^\sigma(u)| + \int_{\partial\Omega} T_k^\sigma(u) d\mathcal{H}^{N-1} \leq \int_{\Omega} h(u) f T_k^\sigma(u) dx \stackrel{(4.21)}{=} - \int_{\Omega} (T_k^\sigma(u))^* \operatorname{div} z.$$

Now an application of Green's identity (2.5) takes to

$$\begin{aligned} \int_{\Omega} |DT_k^\sigma(u)| + \int_{\partial\Omega} T_k^\sigma(u) d\mathcal{H}^{N-1} &\leq \int_{\Omega} (z, DT_k^\sigma(u)) - \int_{\partial\Omega} [T_k^\sigma(u)z, \nu] d\mathcal{H}^{N-1} \\ &= \int_{\Omega} |DT_k^\sigma(u)| - \int_{\partial\Omega} T_k^\sigma(u)[z, \nu] d\mathcal{H}^{N-1}, \end{aligned} \quad (4.26)$$

where we have used (4.23) and the fact that $z \in \mathcal{DM}^\infty(\Omega)$. In particular (4.26) implies that

$$T_k^\sigma(u)(1 + [z, \nu]) = 0 \quad \mathcal{H}^{N-1}\text{-a.e. on } \partial\Omega,$$

which means $[z, \nu](x) = -1$ or $T_k^\sigma(u(x)) = 0$ for \mathcal{H}^{N-1} -a.e. $x \in \partial\Omega$. Now, let $x \in \partial\Omega$ such that $T_k^\sigma(u(x)) = 0$, then

$$\lim_{\epsilon \rightarrow 0^+} \int_{\Omega \cap B(x, \epsilon)} T_k^\sigma(u(y)) dy = 0.$$

If $\sigma > 1$, by means of the Hölder inequality one gets,

$$\int_{\Omega \cap B(x, \epsilon)} T_k(u(y)) dy \leq \left(\int_{\Omega \cap B(x, \epsilon)} T_k^\sigma(u(y)) dy \right)^{\frac{1}{\sigma}} \xrightarrow{\epsilon \rightarrow 0^+} 0,$$

which concludes the proof. \square

Proof of Theorem 3.3. Let u_p be a solution to (4.1). Then by Corollary 4.5 there exists u finite almost everywhere such that u_p converges almost everywhere to u as $p \rightarrow 1^+$ and such that $T_k(u) \in BV_{\text{loc}}(\Omega)$ for any $k > 0$. Applying now Lemma 4.6 there exists a vector field $z \in \mathcal{DM}^\infty(\Omega)$ with $\|z\|_{L^\infty(\Omega)^N} \leq 1$ such that (3.2) and (3.3) hold. The same lemma gives that $h(u)f$ is integrable. Moreover condition (3.4) is proved in Lemma 4.7, and this concludes the proof of Theorem 3.3. \square

Proof of Proposition 3.5. Let u be a solution to (3.1), so that there exists $z \in L^\infty(\Omega)^N$ such that $\|z\|_{L^\infty(\Omega)^N} \leq 1$ and

$$-\operatorname{div} z = h(u)f \quad \text{in } \mathcal{D}'(\Omega). \quad (4.27)$$

Consider $v \in W_0^{1,1}(\Omega)$, multiply (4.27) by $|T_k(v)|$ and apply Green's formula, then

$$\int_{\Omega} h(u)f |T_k(v)| dx = \int_{\Omega} z \cdot \nabla |T_k(v)| dx \leq \|z\|_{L^\infty(\Omega)^N} \int_{\Omega} |\nabla |T_k(v)|| dx \leq \int_{\Omega} |\nabla v| dx.$$

Letting k go to infinity, we get

$$\left| \int_{\Omega} h(u)f v dx \right| \leq \int_{\Omega} h(u)f |v| dx \leq \int_{\Omega} |\nabla v| dx.$$

Since it holds for every $v \in W_0^{1,1}(\Omega)$, it follows that $h(u)f \in W^{-1,\infty}(\Omega)$ and its norm is less than 1. \square

Proof of Proposition 3.6. The solution to (3.1) given by Theorem 3.3 is constructed through u_p solutions to (3.1). Hence, having $T_k^\sigma(u) \in BV(\Omega)$ for any $k > 0$ is consequence of Lemma 4.4. Finally, recalling that $h(u)f \in L_{\text{loc}}^1(\Omega)$, we infer that if $h(0) = \infty$, then $u > 0$ in Ω since $f > 0$ in Ω . \square

4.4. Proof of the uniqueness result. In this subsection we prove the uniqueness result stated in Theorem 3.4.

Proof of Theorem 3.4. Let u be a solution to (3.1), as $\operatorname{div} z \in L^1(\Omega)$ one easily gets from (4.27)

$$-\int_{\Omega} v \operatorname{div} z \, dx = \int_{\Omega} h(u) f v \, dx, \quad (4.28)$$

for all $v \in BV(\Omega) \cap L^\infty(\Omega)$. Suppose that u_1 and u_2 are two solutions to (3.1). Their associated vector fields are, respectively, z_1 and z_2 . At this point we take $v = T_k^\sigma(u_1) - T_k^\sigma(u_2)$ ($k > 0$) in the difference among the variational formulations (4.28) solved, resp., by u_1 and u_2 .

This takes us to

$$\begin{aligned} & \int_{\Omega} (z_1, DT_k^\sigma(u_1)) - \int_{\Omega} (z_1, DT_k^\sigma(u_2)) - \int_{\Omega} (z_2, DT_k^\sigma(u_1)) + \int_{\Omega} (z_2, DT_k^\sigma(u_2)) \\ & - \int_{\partial\Omega} (T_k^\sigma(u_1) - T_k^\sigma(u_2)) [z_1, \nu] \, d\mathcal{H}^{N-1} + \int_{\partial\Omega} (T_k^\sigma(u_1) - T_k^\sigma(u_2)) [z_2, \nu] \, d\mathcal{H}^{N-1} \\ & = \int_{\Omega} (h(u_1) - h(u_2)) f (T_k^\sigma(u_1) - T_k^\sigma(u_2)) \, dx, \end{aligned}$$

where we have exploited (2.5). Moreover, recalling (3.3), one can use again (4.24) deducing that

$$(z_i, DT_k^\sigma(u_i)) = |DT_k^\sigma(u_i)| \quad \text{as measures in } \Omega, \quad i = 1, 2,$$

and recalling (3.4), one yields to

$$\begin{aligned} & \int_{\Omega} |DT_k^\sigma(u_1)| - \int_{\Omega} (z_1, DT_k^\sigma(u_2)) - \int_{\Omega} (z_2, DT_k^\sigma(u_1)) + \int_{\Omega} |DT_k^\sigma(u_2)| \\ & + \int_{\partial\Omega} (T_k^\sigma(u_1) + T_k^\sigma(u_1)) [z_2, \nu] \, d\mathcal{H}^{N-1} + \int_{\partial\Omega} (T_k^\sigma(u_2) + T_k^\sigma(u_2)) [z_1, \nu] \, d\mathcal{H}^{N-1} \\ & \leq \int_{\Omega} (h(u_1) - h(u_2)) f (T_k^\sigma(u_1) - T_k^\sigma(u_2)) \, dx \leq 0. \end{aligned}$$

From the previous inequalities and from the facts that $\|z_i\|_{L^\infty(\Omega)^N} \leq 1$, (2.3), and that $[z_i, \nu] \in [-1, 1]$ for $i = 1, 2$, one deduces that

$$\int_{\Omega} (h(u_1) - h(u_2)) f (T_k^\sigma(u_1) - T_k^\sigma(u_2)) \, dx = 0,$$

which, since h is decreasing and $f > 0$, gives $T_k^\sigma(u_1) = T_k^\sigma(u_2)$ a.e. in Ω for every $k > 0$. In particular $u_1 = u_2$ a.e. in Ω and the proof is concluded. \square

5. THE CASE OF A NONNEGATIVE f

In this section we treat the case of a purely nonnegative datum f . The main point here is that one is not able to show the strict positivity of the solution as a consequence of (4.15). As we will see in Section 6.3 below, this is not a merely technical issue but a structural one and it is closely related to a non-uniqueness phenomenon.

We define the following function

$$\psi(s) = \begin{cases} 1 & \text{if } h(0) < \infty, \\ \chi_{\{s>0\}} & \text{if } h(0) = \infty. \end{cases}$$

Now we are ready to give a meaning to a solution of (3.1) in the case of a nonnegative datum.

Definition 5.1. A nonnegative function u having $T_k(u) \in BV_{\text{loc}}(\Omega)$, $T_k^\sigma(u) \in BV(\Omega)$ for any $k > 0$ and $\chi_{\{u>0\}} \in BV_{\text{loc}}(\Omega)$ is a solution to problem (3.1) if $h(u)f \in L_{\text{loc}}^1(\Omega)$, there exists $z \in \mathcal{DM}_{\text{loc}}^\infty(\Omega)$

such that $\|z\|_{L^\infty(\Omega)^N} \leq 1$ and

$$-\psi^*(u) \operatorname{div} z = h(u)f \quad \text{as measures in } \Omega, \quad (5.1)$$

$$(z, DT_k(u)) = |DT_k(u)| \quad \text{as measures in } \Omega \text{ for any } k > 0, \quad (5.2)$$

$$T_k^\sigma(u(x)) + [T_k^\sigma(u)z, \nu](x) = 0 \quad \text{for } \mathcal{H}^{N-1}\text{-a.e. } x \in \partial\Omega \text{ and for any } k > 0. \quad (5.3)$$

Remark 5.2. It should be noted that Definition 5.1 (needed for purely nonnegative data f) is nothing but a direct extension of Definition 3.1. Indeed if $f > 0$ in Ω and $h(0) = \infty$ (the finite case being simpler) the request $h(u)f \in L_{\text{loc}}^1(\Omega)$ gives that $u > 0$ in Ω and condition (5.1) becomes $-\operatorname{div} z = h(u)f$. This allows us to apply Lemma 5.3 of [12] in order to deduce that $z \in \mathcal{DM}^\infty(\Omega)$, and then, recalling (2.4), one has that

$$[T_k^\sigma(u)z, \nu] = T_k^\sigma(u)[z, \nu],$$

which implies that (5.3) becomes

$$T_k^\sigma(u(x))(1 + [z, \nu](x)) = 0 \quad \text{for } \mathcal{H}^{N-1}\text{-a.e. } x \in \partial\Omega \text{ and for any } k > 0.$$

This identity implies (3.4).

We state and prove the existence theorem for this case:

Theorem 5.3. *Let $0 \leq f \in L^1(\Omega)$ and let h satisfy (h1) and (h2) with $h(\infty) = 0$. Then there exists a solution u to problem (3.1) in the sense of Definition 5.1.*

Proof. We only sketch the proof, underlining the main differences with respect to the one of Theorem 3.3. We consider u_p solutions to (4.1) and we apply Corollary 4.5 in order to deduce the existence of a function u (finite almost everywhere) such that, up to subsequences, u_p converges almost everywhere to u as $p \rightarrow 1^+$. Moreover, one has $T_k^\sigma(u) \in BV(\Omega)$ and $T_k(u) \in BV_{\text{loc}}(\Omega)$ for any $k > 0$. The existence of a vector field $z \in \mathcal{DM}_{\text{loc}}^\infty(\Omega)$ with $\|z\|_{L^\infty(\Omega)^N} \leq 1$ (weak limit in $L^q(\Omega)^N$ of $(|\nabla u_p|^{p-2} \nabla u_p)$ for every $1 \leq q < \infty$) follows exactly as in case $f > 0$. Furthermore, analogously to the proof of Lemma 4.6, it can be proven that $h(u)f \in L_{\text{loc}}^1(\Omega)$ and that (5.2) holds. The proof of (5.3) follows as in Lemma 4.7 up to:

$$\begin{aligned} \int_{\Omega} |DT_k^\sigma(u)| + \int_{\partial\Omega} T_k^\sigma(u) d\mathcal{H}^{N-1} &\leq \int_{\Omega} (z, DT_k^\sigma(u)) - \int_{\partial\Omega} [T_k^\sigma(u)z, \nu] d\mathcal{H}^{N-1} \\ &\leq \int_{\Omega} |DT_k^\sigma(u)| - \int_{\partial\Omega} [T_k^\sigma(u)z, \nu] d\mathcal{H}^{N-1}, \end{aligned}$$

from which (5.3) follows.

Now, we need to show that (5.1) holds and that $\chi_{\{u>0\}} \in BV_{\text{loc}}(\Omega)$. If $h(0) < \infty$ the proof of (5.1) is trivial; hence we assume that $h(0) = \infty$ and we test (4.1) with $R_\delta(u_p)\varphi$ where $R_\delta(s) := 1 - V_\delta(s)$ (V_δ is defined in (2.1)) and $0 \leq \varphi \in C_c^1(\Omega)$. After an application of the Young inequality, one gains

$$\int_{\Omega} |\nabla R_\delta(u_p)| \varphi dx + \int_{\Omega} |\nabla u_p|^{p-2} \nabla u_p \cdot \nabla \varphi R_\delta(u_p) dx \leq \frac{p-1}{p} \int_{\Omega} \varphi dx + \int_{\Omega} h_p(u_p) f_p R_\delta(u_p) \varphi dx,$$

and we intend to pass to the limit first as $p \rightarrow 1^+$, and then as $\delta \rightarrow 0^+$. First of all we can pass by lower semicontinuity in the first term on the left hand; the second term easily passes to the limit as well. Regarding the right hand side we have that the first term vanishes while the second term passes to the limit via dominated convergence (recall that we are integrating where $u_p \geq \delta$). Hence one has

$$\int_{\Omega} |DR_\delta(u)| \varphi dx + \int_{\Omega} z \cdot \nabla \varphi R_\delta(u) dx \leq \int_{\Omega} h(u)f R_\delta(u) \varphi dx.$$

Since $h(u)f \in L_{\text{loc}}^1(\Omega)$ and $z \in L^\infty(\Omega)^N$ one can pass to the limit the previous inequality as $\delta \rightarrow 0^+$, obtaining

$$\int_{\Omega} |D\chi_{\{u>0\}}| \varphi + \int_{\Omega} z \cdot \nabla \varphi \chi_{\{u>0\}} dx \leq \int_{\Omega} h(u)f \chi_{\{u>0\}} \varphi dx = \int_{\Omega} h(u)f \varphi dx,$$

where the last equality holds since $h(u)f \in L^1_{\text{loc}}(\Omega)$ implies that $\{u = 0\} \subset \{f = 0\}$. Moreover this inequality also provides that $\chi_{\{u>0\}} \in BV_{\text{loc}}(\Omega)$. Therefore, it follows from (2.2) that one has

$$-\int_{\Omega} \chi_{\{u>0\}}^* \varphi \operatorname{div} z \leq \int_{\Omega} h(u) f \varphi \, dx. \quad (5.4)$$

We just need to prove the reverse inequality to (5.4). From the weak formulation of (4.1), taking a nonnegative $\varphi \in C_c^\infty(\Omega)$ and letting $p \rightarrow 1^+$ by the Fatou Lemma one has

$$-\int_{\Omega} \varphi \operatorname{div} z \, dx = \int_{\Omega} z \cdot \nabla \varphi \, dx \geq \int_{\Omega} h(u) f \, dx.$$

In particular, we can take $\varphi = (\chi_{\{u>0\}} * \rho_\epsilon) \phi$ where $0 \leq \phi \in C_c^1(\Omega)$ and ρ_ϵ is again a sequence of standard mollifiers. Thus one can pass to the limit in ϵ , obtaining

$$-\int_{\Omega} \phi \chi_{\{u>0\}}^* \operatorname{div} z \geq \int_{\Omega} h(u) f \chi_{\{u>0\}} \phi \, dx = \int_{\Omega} h(u) f \phi \, dx \quad \forall \phi \in C_c^1(\Omega), \quad \phi \geq 0,$$

which proves (5.1). This concludes the proof. \square

6. REMARKS AND EXAMPLES

6.1. The case of a nonnegative h . We want to show that in case in which h touches zero then one can find a bounded solution even if f is merely integrable. Let us suppose that there exists $\tilde{s} > 0$ such that $h(\tilde{s}) = 0$ and $h(s) > 0$ for all $0 < s < \tilde{s}$. We consider the approximation given by (4.1) where

$$h_p(s) = \begin{cases} T_{\frac{1}{p-1}}(h(s)), & 0 \leq s \leq \tilde{s}, \\ 0, & s > \tilde{s}. \end{cases}$$

and taking $(u_p - T_{\tilde{s}}(u_p))^+$ as a test function in (4.1) we deduce that

$$\int_{\Omega} |\nabla(u_p - T_{\tilde{s}}(u_p))^+|^p \, dx \leq 0,$$

which implies that $u_p \leq \tilde{s}$. Hence, reasoning as in the proof of Theorem 3.3, one can prove that there exists a solution u to problem (3.1) which belongs to $L^\infty(\Omega)$ and satisfies $\|u\|_{L^\infty(\Omega)} \leq \tilde{s}$. It is worth remarking that the solution to this problem may not be unique (see Example 3 below).

6.2. The case $h(\infty) > 0$. Let us now focus instead on the case in which $h(s) > 0$ and

$$h(\infty) = \lim_{s \rightarrow \infty} h(s) > 0.$$

Then, the function h is bounded from below by $m > 0$. Assuming that for some positive data f there exists a solution u with vector field z , it holds

$$-\operatorname{div} z = h(u) f \geq m f \text{ in } \Omega.$$

We use now a test function $v \in W_0^{1,1}(\Omega)$ to obtain

$$\int_{\Omega} |\nabla v| \, dx \geq m \int_{\Omega} f |v| \, dx,$$

which implies

$$\left| \int_{\Omega} f v \, dx \right| \leq \int_{\Omega} f |v| \, dx \leq \frac{1}{m} \int_{\Omega} |\nabla v| \, dx,$$

and one may deduce that, if exists a solution, then $f \in W^{-1,\infty}(\Omega)$ and $\|f\|_{W^{-1,\infty}(\Omega)} \leq \frac{1}{m}$. Moreover, the closer to 0 is function h , the more general conditions over f in order to have a solution.

One may also guess that, suitably weakening the definition, one can get existence of a solution; the proof, as $h(s) \geq m > 0$, being a step-by-step replica of the one in [21]. Only observe that in this case the solution u may be, in general, $+\infty$ on a set of positive measure as well as the vector field z could be unbounded.

6.3. Examples. This section is devoted to show some explicit examples of solutions; we look for radial solutions. Let $\Omega = B_R(0)$ be a ball centered at the origin of radius $R > 0$. Let $h(s) = s^{-\gamma}$, $\gamma > 0$, that is we focus on

$$\begin{cases} -\operatorname{div} \left(\frac{Du}{|Du|} \right) = \frac{f(x)}{u^\gamma} & \text{in } B_R(0), \\ u(x) \geq 0 & \text{in } B_R(0), \\ u(x) = 0 & \text{on } \partial B_R(0), \end{cases} \quad (6.1)$$

for various type of data f . We look for non-increasing radial solutions, i.e., $u(x) = g(|x|)$ with $g'(s) \leq 0$ for $0 \leq s \leq R$. Since solutions have to be nonnegative, the boundary condition becomes either $u|_{\partial B_R(0)} = 0$ or $[z, \nu] = -1$. We remark that if $g'(s) < 0$ in a zone of positive measure, then $z(x) = \frac{Du}{|Du|} = -\frac{x}{|x|}$ and $-\operatorname{div} z = \frac{N-1}{|x|}$.

In the first example we construct a solution for a datum f with summability between 1 and N :

Example 1. We start taking the positive datum $f(x) = \frac{N-1}{|x|^q}$ with $1 < q < N$ in problem (6.1):

$$\begin{cases} -\operatorname{div} \left(\frac{Du}{|Du|} \right) = \frac{N-1}{|x|^q} \frac{1}{u^\gamma(x)} & \text{in } B_R(0), \\ u(x) \geq 0 & \text{in } B_R(0), \\ u(x) = 0 & \text{on } \partial B_R(0). \end{cases}$$

Assuming that solution u is non-constant (which implies that the vector field is given by $z = -x/|x|$) and taking $|x| = r$, then equation becomes

$$\frac{N-1}{r} = -\operatorname{div} z = \frac{N-1}{r^q} \frac{1}{g^\gamma(r)},$$

and therefore, the unique solution to this problem is given by $u(x) = g(|x|) = |x|^{\frac{1-q}{\gamma}}$. Observe that u is positive, it has truncation in $BV(B_R(0))$, but is unbounded. Moreover, it holds the equality $(z(x), DT_k(u(x))) = |DT_k(u(x))|$, for any $k > 0$, as measures in $B_R(0)$ and the boundary condition $[z, \nu] = -1$.

Note also that for $q \rightarrow 1^+$ solutions formally tend to a bounded function as the ones found in [12]. Also notice that, if $\gamma = 1$, as $q \rightarrow N^-$ (i.e. the datum goes narrowly outside L^1) then u , in the limit, comes out to $L^{1^+}(B_R(0))$, yielding a sort of *asymptotic optimality* to our result.

Furthermore, assume $R > 1$ and observe that if we formally let $\gamma \rightarrow 0^+$ then the solutions tend to

$$u_0(x) = \begin{cases} +\infty & \text{if } 0 < |x| < 1, \\ 1 & \text{if } |x| = 1, \\ 0 & \text{if } 1 < |x| < R, \end{cases}$$

recovering the blow-up/degenerate phenomenon in [20, Theorem 3.1].


The next example illustrates the case of a generic nonnegative datum f (i.e. the situation of Section 5) and shows that solutions may vanishes on a set of positive measure. Here for simplicity we assume $\gamma = 1$.

Example 2. Consider

$$f(x) = \begin{cases} \frac{N}{\rho} & \text{if } 0 < |x| \leq \rho, \\ 0 & \text{if } \rho < |x| < R, \end{cases}$$

for some $0 < \rho < R$; we define the vector field

$$z(x) = \begin{cases} \frac{-x}{\rho} & \text{if } 0 < |x| \leq \rho, \\ \frac{-x}{|x|^N} \rho^{N-1} & \text{if } \rho < |x| < R, \end{cases}$$

FIGURE 3. The limit $u_0(x)$ as $\gamma \rightarrow 0^+$.

which implies

$$-\operatorname{div} z(x) = \begin{cases} \frac{N}{\rho} & \text{if } 0 < |x| \leq \rho, \\ 0 & \text{if } \rho < |x| < R. \end{cases}$$

Then the solution of (6.1) (with $\gamma = 1$) is given by

$$u(x) = \begin{cases} 1 & \text{if } 0 < |x| \leq \rho, \\ 0 & \text{if } \rho < |x| < R. \end{cases}$$

In fact, it is easy to check that $u \in BV(B_R(0))$, boundary condition holds as well as $(z(x), Du(x)) = |Du(x)|$ since

$$Du(x) = (u^+(x) - u^-(x))v(x)\mathcal{H}^{N-1}\llcorner_{\{|x|=\rho\}} = \frac{-x}{\rho}\mathcal{H}^{N-1}\llcorner_{\{|x|=\rho\}},$$

so that

$$(z(x), Du(x)) = \left(\frac{-x}{\rho}, \frac{-x}{\rho} \right) \mathcal{H}^{N-1}\llcorner_{\{|x|=\rho\}} = |Du(x)|.$$

In this last example we emphasize that the previous degeneracy is not caused by the smoothness of the datum f and, moreover, we provide a non-uniqueness instance.

Example 3. We now consider, in problem (6.1) with $\gamma = 1$, the nonnegative datum

$$f(x) = \begin{cases} \frac{N-1}{|x|^q} & \text{if } 0 < |x| \leq \rho, \\ 0 & \text{if } \rho < |x| < R, \end{cases}$$

with $0 < \rho < R$, $1 < q < N$. If $0 \leq r \leq \rho$, the vector field is given by $z(x) = -x/|x|$ and the solution is $u(x) = 1/|x|^{q-1}$; observe that in this case u itself is $BV(B_R(0))$. Taking the vector field $z(x) = -x|x|^{-N}\rho^{N-1}$ in $\rho < r \leq R$, it is easy to check that the solution is given by $u(x) = \rho^{1-q}$ for $\rho < |x| < R$. Since the boundary condition holds ($[z, v] = (-x/R, x/R) = -1$) as well as the equality $(z(x), Du(x)) = |Du(x)|$ as measures, we conclude that u is indeed a solution of (6.1). Moreover, uniqueness does not hold owing to function

$$v(x) = \begin{cases} \frac{1}{|x|^{q-1}} & \text{if } 0 < |x| < \rho, \\ 0 & \text{if } \rho < |x| < R, \end{cases}$$

is also a solution with the same vector field. Indeed, $v \in BV(B_R(0))$, it is straightforward that boundary condition is satisfied and $(z(x), Dv(x)) = |Dv(x)|$ holds due to

$$\begin{aligned} Dv(x)\llcorner_{\{|x|=\rho\}} &= (v^+(x) - v^-(x))v(x)\mathcal{H}^{N-1}\llcorner_{\{|x|=\rho\}} = \left(\frac{1}{\rho^{q-1}} \right) \left(\frac{-x}{\rho} \right) \mathcal{H}^{N-1}\llcorner_{\{|x|=\rho\}} \\ &= \frac{-x}{\rho^q} \mathcal{H}^{N-1}\llcorner_{\{|x|=\rho\}}, \end{aligned}$$

and so

$$(z(x), Dv(x))_{\mathcal{L}_{\{|x|=\rho\}}} = \frac{1}{\rho^{q-1}} \mathcal{H}^{N-1}_{\{|x|=\rho\}} = |Dv(x)|_{\mathcal{L}_{\{|x|=\rho\}}}.$$

ACKNOWLEDGEMENT

The fourth author is supported by the Spanish Ministerio de Ciencia, Innovación y Universidades and FEDER, under project PGC2018–094775–B–I00.

REFERENCES

- [1] L. Ambrosio, N. Fusco and D. Pallara, *Functions of Bounded Variation and Free Discontinuity Problems*, Oxford Mathematical Monographs, 2000.
- [2] F. Andreu, C. Ballester, V. Caselles and J.M. Mazón, The Dirichlet problem for the total variation flow, *J. Funct. Anal.* 180 (2) (2001) 347–403.
- [3] F. Andreu, V. Caselles and J. M. Mazón, *Parabolic quasilinear equations minimizing linear growth functionals*, Progress in Mathematics, 223, Birkhäuser Verlag, Basel, 2004
- [4] G. Anzellotti, Pairings between measures and bounded functions and compensated compactness, *Ann. Mat. Pura Appl.* 135 (4) (1983) 293–318.
- [5] M. Bertalmio, V. Caselles, B. Rougé and A. Solé: TV based image restoration with local constraints. Special issue in honor of the sixtieth birthday of Stanley Osher, *J. Sci. Comput.* 19 (1-3) (2003) 95–122.
- [6] L. Boccardo and L. Orsina, Semilinear elliptic equations with singular nonlinearities, *Calc. Var. Partial Differential Equations* 37 (2010) 363–380.
- [7] V. Caselles, On the entropy conditions for some flux limited diffusion equations, *Journal of Differential Equations* 250 (2011) 3311–3348.
- [8] K.C. Chang, The spectrum of the 1–Laplace operator. *Commun. Contemp. Math.* 11 (2009), no. 5, 865–894.
- [9] G. Q. Chen and H. Frid, Divergence-measure fields and hyperbolic conservation laws, *Arch. Ration. Mech. Anal.* 147 (2) (1999) 89–118.
- [10] M. Cicalese and C. Trombetti, Asymptotic behaviour of solutions to p –Laplacian equation, *Asymptot. Anal.* 35 (1) (2003) 27–40.
- [11] G. Crasta and V. De Cicco, Anzellotti’s pairing theory and the Gauss–Green theorem, *Advances in Mathematics* 343 (2019) 935–970.
- [12] V. De Cicco, D. Giachetti, F. Oliva and F. Petitta, The Dirichlet problem for singular elliptic equations with general nonlinearities, *Calculus of Variations and Partial Differential Equations* 58 (4) (2019) 129.
- [13] V. De Cicco, D. Giachetti and S. Segura de León, Elliptic problems involving the 1–Laplacian and a singular lower order term, *J. Lond. Math. Soc.* (2) 99 (2019) 349–376.
- [14] F. Demengel, On some nonlinear partial differential equations involving the “1”–Laplacian and critical Sobolev exponent, *ESAIM, Control Optim. Calc. Var.* 4 (1999) 667–686.
- [15] G.M. Figueiredo and M.T.O. Pimenta, Nehari method for locally Lipschitz functionals with examples in problems in the space of bounded variation functions, *NoDEA, Nonlinear Differ. Equ. Appl.* 25 (2018) No. 5, Paper No. 47, 18 p.
- [16] B. Kawohl, On a family of torsional creep problems, *J. Reine Angew. Math.* 410 (1990) 1–22.
- [17] B. Kawohl, From p –Laplace to mean curvature operator and related questions, In : *Progress in Partial Differential Equations: the Metz Surveys*, Pitman Res. Notes Math. Ser., Vol. 249, Longman Sci. Tech., Harlow, pp. 40–56, 1991
- [18] B. Kawohl and F. Schuricht, Dirichlet problems for the 1–Laplace operator, including the eigenvalue problem, *Commun. Contemp. Math.* 9 (4) (2007) 515–543.
- [19] J. Leray and J.-L. Lions, Quelques résultats de Višik sur les problèmes elliptiques non linéaires par les méthodes de Minty–Browder, *Bull. Soc. Math. Fr.* 93 (1965) 97–107.
- [20] A. Mercaldo, S. Segura de León and C. Trombetti, On the behaviour of the solutions to p –Laplacian equations as p goes to 1, *Publicacions Matemàtiques* 52 (2) (2008) 377–411.
- [21] A. Mercaldo, S. Segura de León and C. Trombetti, On the solutions to 1–Laplacian equation with L^1 data, *Journal of Functional Analysis* 256 (8) (2009) 2387–2416.
- [22] Y. Meyer, *Oscillating Patterns in Image Processing and Nonlinear Evolution Equations: The Fifteenth Dean Jacqueline B. Lewis Memorial Lectures*, Providence, RI: American Mathematical Society, 2001.
- [23] A. Molino Salas and S. Segura de León, Elliptic equations involving the 1–Laplacian and a subcritical source term, *Nonlinear Anal., Theory Methods Appl., Ser. A, Theory Methods* 168, (2018) 50–66.
- [24] F. Oliva, Regularizing effect of absorption terms in singular problems, *J. Math. Anal. Appl.* 472 (1) (2019) 1136–1166.
- [25] F. Oliva and F. Petitta, Finite and infinite energy solutions of singular elliptic problems: Existence and uniqueness, *J. Differential Equations* 264 (1) (2018) 311–340.

- [26] S. Osher and J. Sethian, Fronts propagating with curvature-dependent speed: algorithms based on Hamilton-Jacobi formulations, *Journal of Computational Physics* 79 (1) (1988) 12–49.
- [27] G. Sapiro, *Geometric partial differential equations and image analysis*, Cambridge University Press, 2001.
- [28] A.C.M. van Rooij and W.H. Schikhof, *A second course on real functions*, Cambridge University Press, 1982.

MARTA LATORRE

MATEMÁTICA APLICADA, CIENCIA E INGENIERÍA DE LOS MATERIALES Y TECNOLOGÍA ELECTRÓNICA, UNIVERSIDAD REY JUAN CARLOS

C/TULIPÁN S/N 28933, MÓSTOLES, SPAIN

E-mail address: marta.latorre@urjc.es

FRANCESCANTONIO OLIVA

DIPARTIMENTO DI SCIENZE DI BASE E APPLICATE PER L' INGEGNERIA, SAPIENZA UNIVERSITÀ DI ROMA

VIA SCARPA 16, 00161 ROMA, ITALY

E-mail address: francesco.oliva@sbai.uniroma1.it

FRANCESCO PETITTA

DIPARTIMENTO DI SCIENZE DI BASE E APPLICATE PER L' INGEGNERIA, SAPIENZA UNIVERSITÀ DI ROMA

VIA SCARPA 16, 00161 ROMA, ITALY

E-mail address: francesco.petitta@sbai.uniroma1.it

SERGIO SEGURA DE LEÓN

DEPARTAMENT D'ANÀLISI MATEMÀTICA, UNIVERSITAT DE VALÈNCIA

DR. MOLINER 50, 46100 BURJASSOT, VALÈNCIA, SPAIN

E-mail address: sergio.segura@uv.es