

## POSITIVE $p$ -SUMMING OPERATORS, VECTOR MEASURES AND TENSOR PRODUCTS

by OSCAR BLASCO

(Received 25th February 1986)

### Introduction

In this paper we shall introduce a certain class of operators from a Banach lattice  $X$  into a Banach space  $B$  (see Definition 1) which is closely related to  $p$ -absolutely summing operators defined by Pietsch [8].

These operators, called positive  $p$ -summing, have already been considered in [9] in the case  $p=1$  (there they are called cone absolutely summing, c.a.s.) and in [1] by the author who found this space to be the space of boundary values of harmonic  $B$ -valued functions in  $h_B^p(D)$ .

Here we shall use these spaces and the space of majorizing operators to characterize the space of bounded  $p$ -variation measures  $V_B^p$  and to endow the tensor product  $L^p \otimes B$  with a norm in order to get  $L^p(B)$  as its completion in this norm.

### Some definitions and previous results

Throughout this paper  $X$  will denote a Banach lattice and  $B$  a Banach space. Given  $1 \leq p \leq \infty$  we shall always write  $p'$  for such a number that  $(1/p) + (1/p') = 1$ .

**Definition 1.** An operator  $T$  belonging to  $L(X, B)$  is called *positive  $p$ -summing* ( $1 \leq p < \infty$ ) if there exists a constant  $C > 0$  such that for all *positive* elements  $x_1, x_2, \dots, x_n$  in  $X$  we have

$$\left( \sum_{i=1}^n \|Tx_i\|_B^p \right)^{1/p} \leq C \cdot \sup_{\|\xi\|_{X^*} \leq 1} \left( \sum_{i=1}^n |\langle \xi, x_i \rangle|^p \right)^{1/p}. \quad (1)$$

We shall denote by  $\Lambda_p(X, B)$  the space of such operators and the infimum of the constants will be the norm on it.

A duality argument allows us to write the following equivalent formulation of (1):

$$\left( \sum_{i=1}^n \|Tx_i\|_B^p \right)^{1/p} \leq C \cdot \sup \left\{ \left\| \sum_{i=1}^n \alpha_i \cdot x_i \right\|_X : \sum_{i=1}^n \alpha_i^{p'} \leq 1, \alpha_i \geq 0 \right\}. \quad (1')$$

Obviously the space of  $p$ -absolutely summing operators  $\Pi_p(X, B)$  is included in  $\Lambda_p(X, B)$  and the same techniques as for  $p$ -absolutely summing operators lead us to see

that for  $p \leq q$ ,  $\Lambda_p(X, B) \subseteq \Lambda_q(X, B)$  and

$$|T|_{\Lambda_q} \leq |T|_{\Lambda_p} \quad \text{for all } T \text{ in } \Lambda_p(X, B). \quad (2)$$

**Definition 2** (see [9]). An operator  $T$  belonging to  $L(B, X)$  is called *majorizing* if there exists a constant  $C > 0$  such that for every  $x_1, x_2, \dots, x_n$  in  $B$

$$\left\| \sup_{1 \leq i \leq n} |Tx_i| \right\|_X \leq C \cdot \sup_{1 \leq i \leq n} \|x_i\|_B. \quad (3)$$

We shall denote by  $M(B, X)$  the space of such operators and we shall set the following norm on it:

$$|T|_m = \sup \left\{ \left\| \sup_{1 \leq i \leq n} |Tx_i| \right\|_X : \{x_i\} \in B, \|x_i\|_B \leq 1 \right\}.$$

If we consider  $A \otimes B$  as a subspace of  $L(A^*, B)$ , that is  $u = \sum_{i=1}^n a_i \otimes b_i$  represents the operator  $T_u$  defined by  $T_u(\xi) = \sum_{i=1}^n \langle \xi, a_i \rangle \cdot b_i$ , then it is easy to see that  $A \otimes B$  is included in  $\Lambda_p(A^*, B)$  and  $M(A^*, B)$ . Let us denote by  $A \hat{\otimes}_p B$  and  $A \check{\otimes}_m B$  the completion of the space  $A \otimes B$  endowed with the norms induced by  $\Lambda_p(A^*, B)$  and  $M(A^*, B)$  respectively.

#### Applications to tensor products and vector measures

Let  $(\Omega, \mathcal{B}, \mu)$  be a finite measure space and  $1 \leq p < \infty$ . We shall denote by  $L^p(\mu, B)$  the space of measurable functions such that  $\|f\|_p = (\int_{\Omega} \|f(t)\|^p d\mu)^{1/p} < +\infty$ .

The following result can be found in [9].

$$L^p(\mu) \hat{\otimes}_1 B = L^p(\mu, B) \quad 1 \leq p < \infty. \quad (4)$$

This fact can be extended in the following way:

**Theorem 1.** Let  $1 \leq p < \infty$ , then for all  $1 \leq r \leq p$

$$L^p(\mu) \hat{\otimes}_r B = L^p(\mu, B).$$

**Proof.** Let  $1 \leq r \leq p$ . Since simple functions are dense in  $L^p(\mu, B)$ , it suffices to show that for each  $s = \sum_{i=1}^n x_i \cdot \chi_{E_i}$  we have that the operator  $T_s(\psi) = \int_{\Omega} s(t) \cdot \psi(t) d\mu(t)$  satisfies  $|T_s|_{\Lambda_r} = \|s\|_p$ .

Since  $\|s\|_p = |T_s|_{\Lambda_1}$  and  $|T_s|_{\Lambda_p} \leq |T_s|_{\Lambda_r} \leq |T_s|_{\Lambda_1}$ , then it is enough to prove that  $\|s\|_p \leq |T_s|_{\Lambda_p}$ .

$$\|s\|_p = \left( \sum_{i=1}^n \|x_i\|^p \cdot \mu(E_i) \right)^{1/p}$$

$$\begin{aligned}
 &= \left( \sum_{i=1}^n \|T_s(\mu(E_i)^{-1} \cdot \chi_{E_i})\|^p \cdot \mu(E_i) \right)^{1/p} \\
 &= \left( \sum_{i=1}^n \|T_s(\mu(E_i)^{-1/p'} \cdot \chi_{E_i})\|^p \right)^{1/p} \\
 &\leq |T_s|_{\Lambda_p} \cdot \sup \left\{ \left\| \sum_{i=1}^n \alpha_i \cdot \mu(E_i)^{-1/p'} \cdot \chi_{E_i} \right\|_{L^{p'}} \mid \sum_{i=1}^n \alpha_i^{p'} \leq 1, \alpha_i \geq 0 \right\} \\
 &= |T_s|_{\Lambda_p}. \quad \square
 \end{aligned}$$

We can give another representation of  $\Lambda_p(L^p(\mu), B)$  in terms of vector measures. Let us recall a space of  $B$ -valued measures, introduced by Bochner [2] in the scalar-valued case, which is a good substitute for  $L^p(\mu, B)$  in several cases, for example for the duality  $(L^p(\mu, B))^* = V_B^{p'}$  or for boundary values of functions in  $h_B^p(D)$  [1].

**Definition 3.** A finitely additive vector measure  $G: \mathcal{B} \rightarrow B$  is said to have bounded  $p$ -variation if

$$|G|_p = \sup_{\pi} \left\{ \left( \sum_{E \in \pi} \frac{\|G(E)\|^p}{\mu(E)^{p-1}} \right)^{1/p} \right\} < +\infty \quad (1 < p < \infty) \tag{5}$$

where the "sup" is taken over all finite partitions of  $\Omega$  and

$$|G|_{\infty} = \sup \left\{ \frac{\|G(E)\|}{\mu(E)}, E \in \mathcal{B} \right\} < +\infty \quad (p = \infty). \tag{5'}$$

We shall denote by  $V_B^p$  the space of such measures and its norm is given by (5) or (5') provided  $1 < p < \infty$  or  $p = \infty$ .

Let us recall some properties of this space.

- (a) Every measure in  $V_B^p$  is countably additive,  $\mu$ -continuous and with bounded variation.
- (b)  $L^p(\mu, B)$  is isometrically embedded in  $V_B^p$ .

Dinculeanu [4] characterized the space  $V_B^p$  in terms of  $\mathcal{L}(L^p(\mu), B)$ , the space of operators in  $L(L^p(\mu), B)$  such that

$$\|T\|_p = \sup \left\{ \sum_{i=1}^n |\alpha_i| \cdot \|T(\chi_{E_i})\|_B \mid \sum_{i=1}^n \alpha_i \cdot \chi_{E_i} \Big\|_{L^{p'}} \leq 1 \right\} < +\infty.$$

The author proved in [1] that  $\mathcal{L}(L^p(\mu), B) = \Lambda_p(L^p(\mu), B)$ , hence we have the following:

**Theorem 2.** For  $1 < p \leq \infty$ ,  $\Lambda_p(L^p(\mu), B) = V_B^p$ .

Now we shall characterize  $V_B^p$  by means of the space of certain majorizing operators.

**Theorem 3.** For  $1 < p < \infty$ ,  $M(B, L^p(\mu)) = V_{B^*}^p$ .

**Proof.** Let  $G$  be a measure of  $V_{B^*}^p$  and take  $x \in B$  with  $\|x\|_B = 1$ . Consider now the measure  $G_x(E) = \langle G(E), x \rangle$  for all measurable set  $E$  and the positive measure  $|G|$ . Both measures are countably additive,  $\mu$ -continuous and with bounded variation. So, by the Radon-Nikodým theorem, there exist  $f_x$  and  $g \geq 0$  in  $L^1(\mu)$  such that

$$G_x(E) = \int_E f_x(t) d\mu(t) \quad \text{for all } E \in \mathcal{B}, \quad (6)$$

$$|G|(E) = \int_E g(t) d\mu(t) \quad \text{for all } E \in \mathcal{B}. \quad (7)$$

It is not difficult to show, since  $G$  belongs to  $V_{B^*}^p$ , that  $f_x$  and  $g$  belong to  $L^p(\mu)$  and moreover  $\|g\|_p = |G|_p$  (see the argument in [1, Proposition 3]).

Due to (6) and (7) we have that

$$|G_x|(E) = \int_E |f_x(t)| d\mu(t) \leq |G|(E) = \int_E g(t) d\mu(t)$$

and from this we obtain

$$|f_x(t)| \leq |g(t)| \quad \mu\text{-a.e.} \quad (8)$$

Let us define  $T: B \rightarrow L^p(\mu)$

$$y \mapsto T(y) = \|y\|_B \cdot f_{y/\|y\|_B}.$$

From (8) it is easy to show that  $T \in M(B, L^p(\mu))$ .

Indeed, if  $x_1, x_2, \dots, x_n$  belong to  $B$  and  $\|x_i\|_B = 1$  then

$$\left\| \sup_{1 \leq i \leq n} |Tx_i| \right\|_{L^p} \leq \|g\|_p = |G|_p.$$

Conversely, given  $T$  in  $M(B, L^p(\mu))$  and denoting by  $f_x$  the function  $Tx$ , we can define the measure  $G: \mathcal{B} \rightarrow B^*$  by

$$\langle G(E), x \rangle = \int_E f_x(t) d\mu(t). \quad (9)$$

Now, let  $\pi$  be a partition of  $\Omega$ . Given  $\varepsilon > 0$ , for each  $E \in \pi$  there exists  $b_E \in B$  with  $\|b_E\|_B = 1$  such that

$$\mu(E)^{-1/p'} \cdot \|G(E)\| \leq \langle \mu(E)^{-1/p'} \cdot G(E), b_E \rangle + \varepsilon/n^{1/p}. \quad (10)$$

From (10) the triangle inequality in  $\ell^p$  implies

$$\left( \sum_{E \in \pi} (\mu(E)^{-1/p'} \cdot \|G(E)\|)^p \right)^{1/p} = \left( \sum_{E \in \pi} |\langle \mu(E)^{-1/p'} \cdot G(E), b_E \rangle|^p \right)^{1/p} + \varepsilon.$$

Now by using (9) we can write

$$\begin{aligned} \left( \sum_{E \in \pi} \frac{\|G(E)\|^p}{\mu(E)^{p-1}} \right)^{1/p} &\leq \left( \sum_{E \in \pi} \left( \mu(E)^{-1/p'} \cdot \left| \int_E f_{b_E}(t) d\mu(t) \right| \right)^p \right)^{1/p} + \varepsilon \\ &= \sup_{\sum \alpha_E^p = 1} \left\{ \sum_{E \in \pi} \int_E |f_{b_E}(t)| \cdot \alpha_E \cdot \mu(E)^{-1/p'} \cdot d\mu(t) \right\} + \varepsilon \\ &\leq \sup_{\sum \alpha_E^p = 1} \left\{ \int_{\Omega} \left( \sup_{E \in \pi} |f_{b_E}(t)| \right) \left( \sum_{E \in \pi} \alpha_E \cdot \mu(E)^{-1/p'} \cdot \chi_E(t) \right) d\mu(t) \right\} + \varepsilon \\ &\leq \left\| \sup_{E \in \pi} |T(b_E)| \right\|_{L^p} \cdot \sup_{\sum \alpha_E^p = 1} \left\| \sum_{E \in \pi} \alpha_E \cdot \mu(E)^{-1/p'} \cdot \chi_E \right\|_{L^{p'}} + \varepsilon \\ &\leq \|T\|_m + \varepsilon. \end{aligned}$$

Taking  $\varepsilon$  arbitrarily small and the "sup" over the partitions we obtain  $\|G\|_p \leq \|T\|_m$ , completing the proof. □

This theorem allows us to prove the following result of [5].

**Corollary.**  $B \otimes_m L^p(\mu) = L^p(\mu, B)$  for each  $1 < p < \infty$ .

**Proof.** Given a simple function  $s = \sum_{i=1}^n x_i \cdot \chi_{E_i}$  where  $x_i$  belongs to  $B$ , we notice that  $s$  clearly belongs to  $L^p(\mu, B^{**})$  and therefore the measure  $G_s(E) = \int_E s(t) d\mu(t)$  belongs to  $V_{B^{**}}^p = M(B^*, L^p(\mu))$ . So, denoting by  $T_s$  the operator associated with  $s$  we have  $\|s\|_p = \|G_s\|_p = \|T_s\|_m$ . Finally the density of simple functions in the space  $L^p(\mu, B)$  gives us the corollary. □

REFERENCES

1. O. BLASCO, Boundary values of vector valued harmonic functions considered as operators, *Studia Math.* **86** (1987), 19-33.
2. S. BOCHNER, Additive set functions on groups, *Ann. of Math.* **40** (1939), 769-799.
3. J. DIESTEL and J. J. UHL, *Vector Measures* (Amer. Math. Soc. Mathematical Surveys **15**, (1977)).
4. N. DINCULEANU, *Vector Measures* (Pergamon Press, New York, 1967).
5. S. HEINRICH, M. J. NIELSEN and G. OLSEN, Order bounded operators and tensor products of Banach lattices, *Math. Scand.* **49** (1981), 99-127.
6. S. LEADER, The theory of  $L^p$ -spaces for finitely additives set functions, *Ann. of Math.* (2) **58** (1953), 528-543.
7. J. LINDENSTRAUSS and L. TZAFIRI, *Classical Banach Spaces, Vols. I and II* (Springer-Verlag, Berlin, 1979).

8. A. PIETSCH, Absolut  $p$ -summierende Abbildungen in normierten Räumen, *Studia Math.* **28** (1967), 333–353.

9. H. H. SCHAEFFER, *Banach Lattices and Positive Operators* (Springer-Verlag, Berlin, 1974).

DPTO. TEORÍA DE FUNCIONES  
FACULTAD DE CIENCIAS  
50009-ZARAGOZA  
SPAIN